

Irritable bowel syndrome in the general population: epidemiology, comorbidity, and societal costs

Markku Hillilä



Helsinki 2010

Division of Gastroenterology, Department of Medicine
Helsinki University Central Hospital
Helsinki, Finland

**Irritable bowel syndrome in the general
population: epidemiology, comorbidity, and
societal costs**

Markku Hillilä

ACADEMIC DISSERTATION

To be presented, with the permission of the Medical Faculty of the University of Helsinki,
for public examination in Lecture Hall 3, Biomedicum Helsinki, Haartmaninkatu 8,
on March 12th, 2010, at 12 noon.

Helsinki 2010

Supervisor

Docent Martti Färkkilä, MD, PhD
Division of Gastroenterology, Department of Medicine
Helsinki University Central Hospital
Helsinki, Finland

Reviewers

Professor Lars Agreus
Centre for Family and Community Medicine
Department of Neurobiology, Care Sciences and Society
Karolinska Institutet
Stockholm, Sweden

Docent Rauli Leino, MD, PhD
Department of Medicine
Turku University Central Hospital
Turku, Finland

Opponent

Docent Markku Heikkinen, MD, PhD
Department of Medicine
Unit of Gastroenterology
Kuopio University Hospital
Kuopio, Finland

ISBN 978-952-92-6896-2 (paperback)
ISBN 978-952-10-6100-4 (PDF)
<http://ethesis.helsinki.fi>

Helsinki University Print
Helsinki 2010

To Anu, Petri, and Taru

Contents

List of original publications	8
Abbreviations	9
Abstract	10
1. INTRODUCTION	12
2. REVIEW OF THE LITERATURE	13
Definition, natural history, and prognosis of irritable bowel syndrome (IBS)	13
Pathogenesis	13
Diagnostic criteria	16
IBS subtypes	17
Epidemiology	19
Prevalence	19
Sociodemographic factors	19
Comorbidity in IBS	21
Psychiatric comorbidity	21
Somatic comorbidity	22
Gastrointestinal comorbidity	22
Extraintestinal comorbidity	23
Treatment options	23
Pharmacological treatment	24
Psychological therapies	25
Healthcare use	26
Abnormal illness behaviour	26
Economic impact	26
Direct costs	27

Indirect costs	28
3. AIMS OF THE STUDY	29
4. MATERIALS AND METHODS	30
Sample	30
Questionnaires	30
Questionnaire I (Appendix A)	30
Pilot study	32
Questionnaire II (Appendix B)	32
Data collection	33
Control groups	33
Statistical methods	34
Ethics	35
5. RESULTS	36
Prevalence of IBS (I)	38
Overlap of IBS criteria (I)	39
GI symptom severity (I)	41
Prevalence of depressive symptoms (III)	41
Comorbidity in IBS (I,II)	41
Healthcare use (I,II,IV)	42
Predictors of healthcare seeking (II, III)	43
Depressive symptoms and health care seeking for GI reasons	45
Societal costs (IV)	45
6. DISCUSSION	47
Methodological aspects	47
Comparison of IBS criteria	48
Depression and anxiety	49

Comorbidity	50
Healthcare seeking	50
Predictors of GI consultations	51
Predictors of non-GI consultations	51
Societal costs	51
SUMMARY AND CONCLUSIONS	54
Acknowledgements	55
References	56
Appendices	
A	Questionnaire I
B	Questionnaire II

List of original publications

This thesis is based on the following publications, which are referred to in the text by their Roman numerals I-IV:

- I Hillilä MT, Färkkilä MA. Prevalence of irritable bowel syndrome according to different diagnostic criteria in a non-selected adult population. *Alimentary Pharmacology & Therapeutics* 2004; 20: 339-345.
- II Hillilä MT, Siivola MT, Färkkilä MA. Comorbidity and use of health-care services among IBS sufferers. *Scandinavian Journal of Gastroenterology* 2007;(42)7:799-806.
- III Hillilä MT, Hämäläinen J, Heikkinen ME, Färkkilä MA. Gastrointestinal complaints among subjects with depressive symptoms in the general population. *Alimentary Pharmacology & Therapeutics* 2008; 28: 648–654.
- IV Hillilä MT, Färkkilä NJ, Färkkilä MA. Societal costs for irritable bowel syndrome – a population based study. *Scandinavian Journal of Gastroenterology*, in press.

The original publications are reprinted with the kind permission of the copyright holders.

Abbreviations

BDI-SF	Beck Depression Inventory Short Form
CI	Confidence interval
CNS	Central nervous system
ENS	Enteric nervous system
FD	Functional dyspepsia
fMRI	Functional magnetic resonance imaging
GERD	Gastro-oesophageal reflux disease
GI	Gastrointestinal
IBS	Irritable bowel syndrome
IBS-C	IBS with constipation
IBS-D	IBS with diarrhoea
IBS-M	Mixed IBS
IgE	Immunoglobulin E
κ	Cohen's kappa value
M	Million
NNT	Number needed to treat
PAN	Primary afferent neuron
PEG	Polyethylene glycol
PET	Proton emission tomography
QoL	Quality of life
R-BDI-SF	Raitasalo-Beck Depression Inventory Short Form
SERT	Serotonin transporter
SSRI	Serotonin reuptake inhibitors
TCA	Tricyclic antidepressant
5-HT	5-hydroxytryptamine

Abstract

Background: Irritable bowel syndrome (IBS) is a functional gastrointestinal (GI) disorder characterised by abdominal pain and abnormal bowel function. The diagnosis of IBS is based on symptom description as no organic, biochemical or structural abnormalities are present. In epidemiologic studies, the prevalence of IBS ranges between 3 and 25%, usually with an over-representation of the female gender. The wide variation in prevalence estimates is partly due to various IBS definitions and diagnostic criteria. The most frequently used symptom based criteria include Manning, Rome I, and Rome II criteria. Manning criteria generally give a higher prevalence estimate than Rome II criteria.

IBS is associated with a high rate of psychological and somatic comorbid conditions, such as depression, anxiety, dyspepsia, headache, lower back pain, and fibromyalgia. Moreover, IBS is associated with a high rate of health care consumption for both GI and non-GI reasons. IBS sufferers miss work for illness almost three times as often as controls. Therefore, IBS incurs significant direct and indirect health care costs for society.

The aim of this study was to assess the prevalence of IBS among subjects of working age according to various diagnostic criteria. In addition, we studied the rates of somatic and psychiatric comorbidity and health care consumption for GI and non-GI reasons, as well as predictors of health care seeking. Finally, we evaluated the societal costs of IBS.

Methods: The study was a two-phase postal survey. In phase I, a questionnaire covering GI symptoms according to Manning 2 (at least two of the six Manning symptoms), Manning 3 (at least three Manning symptoms), Rome I, and Rome II criteria, was mailed to 5 000 randomly selected non-institutionalised Finnish residents aged 18 to 64. The questionnaire also included a Finnish modification of the Beck Depression Inventory Short Form (BDI-SF) and items covering the severity of GI symptoms, rates of headaches, back pain, and dyspeptic symptoms. In addition, health care consumption for GI and non-GI reasons and work absenteeism for GI reasons was inquired. In phase II, a questionnaire covering rates of GI health care visits in primary and secondary care, use of GI medication, and the duration of GI symptoms, was sent to subjects fulfilling IBS criteria according to Manning 2 or Rome II criteria in the phase I questionnaire.

Results: The response rate was 73% and 86% for questionnaires I and II. After adjusting for age and gender, the prevalence of IBS was 15.9% (95% CI 14.7-17.1%), 9.6% (95% CI 8.6-10.6%), 5.6% (95% CI 4.8-6.4%), and 5.1% (95% CI 4.4-5.8%) according to Manning 2, Manning 3, Rome I, and Rome II criteria. Of those meeting Rome II criteria, 97% also met Manning 2 criteria.

The prevalence of depressive symptoms in the general population was 16.7% (95% CI 15.4-17.9%). Of subjects with depressive symptoms, 11.6% met Rome II IBS criteria compared to 3.7% of those with no depressiveness ($p < 0.0001$). In addition, subjects with depressive symptoms reported more physician visits and sick leave days for GI symptoms than those without.

Presence of dyspeptic symptoms, lactose intolerance, headache, back pain, depressive symptoms, anxiety, and insomnia was more common among subjects meeting any IBS

criterion than by controls not meeting the criterion. In addition, presence of asthma or an allergic condition was more common among those fulfilling Manning 2, Manning 3, or Rome II criteria than among controls. Presence of severe or very severe abdominal pain and disturbance of daily activities by abdominal pain were more often reported by subjects meeting either of the Rome criteria than those meeting either of the Manning criteria.

During the previous year, a larger share of subjects meeting any IBS criteria had made GI and non-GI physician visits than controls. In addition, subjects fulfilling either of the Rome criteria more often reported GI physician visits than those fulfilling either of the Manning criteria. Intensity of GI symptoms and presence of dyspeptic symptoms were the strongest predictors of GI consultations, but also headaches, insomnia, and presence of a chronic illness were independent predictors. For non-GI visits, presence of chronic illness, back pain, and depressive symptoms were the strongest predictors. In addition, presence of dyspeptic symptoms and a history of abdominal pain in childhood also predicted non-GI visits.

Annual GI related individual costs were higher in the Rome II group (497€, 95% CI 382-621€) than in the Manning 2 group (295€, 95% CI 246-347€). The nationwide annual GI related expenses were higher by Manning 2 criteria than Rome II criteria (154M€, 95% CI 128-181M€ vs. 82M€, 95% CI 63-103M€). Direct expenses of GI symptoms and non-GI physician visits ranged between 98M€ for Rome II and 230M€ for Manning criteria.

Conclusions: The prevalence of IBS shows substantial variation depending on the criteria applied. Rome II criteria are more restrictive than Manning 2 criteria, and they identify an IBS population with more severe GI symptoms, more frequent health care use, and higher individual health care costs. In the general population, subjects with IBS demonstrate high rates of psychiatric and somatic comorbidity regardless of health care seeking status. Perceived symptom severity rather than psychiatric comorbidity predicts health care seeking for GI symptoms. Depressive symptoms are prevalent in the general population and they are associated with an elevated rate of GI symptoms, physician visits, and work absenteeism for GI symptoms.

IBS imposes a significant impact on national health care expenditures. The direct GI and non-GI costs of IBS are equivalent to up to 5% of outpatient health care and medicine costs in Finland. A more integral approach to IBS by physicians, accounting also for comorbid conditions, may produce a more favourable course in IBS patients and reduce health care expenditures.

1. INTRODUCTION

Irritable bowel syndrome (IBS) is a chronic functional gastrointestinal (GI) disorder characterised by abdominal pain or discomfort and altered bowel habits. The aetiology of IBS is multifactorial with altered visceral sensitivity, altered gastrointestinal motility, and psychosocial factors influencing symptom generation.

The prevalence of IBS varies worldwide, affecting 3 to 25% of the population. Up to 90% of subjects with IBS demonstrate comorbid somatic or psychiatric symptoms. They make more health care visits than the control population without IBS. Work absenteeism due to IBS symptoms is as frequent as that of common cold. Direct and indirect costs of IBS incur considerable societal economic burden.

The definition and diagnostic criteria of IBS has changed over time. In 1978 Manning *et al.* presented the first symptom based criteria for IBS, widely applied later in epidemiological research. In 1989, the Rome criteria were introduced and later modified as Rome I, Rome II, and Rome III criteria.

The wide variation in the prevalence of IBS may reflect true differences between various populations and countries. Differences in IBS definitions and study methodologies, however, hamper reliable comparison between previous studies.

The main aim of this thesis was to assess the prevalence of IBS according to varying diagnostic criteria in a randomly selected population sample and to compare sociodemographic, symptom characteristics, and health care use between subjects meeting varying IBS criteria and a control population. The thesis also aimed to compare rates of comorbid psychiatric and somatic conditions between those meeting IBS criteria and a control population, and evaluate their influence on the use of health care facilities among IBS sufferers. In addition, societal costs of IBS were to be evaluated.

2. REVIEW OF THE LITERATURE

Definition, natural history, and prognosis of irritable bowel syndrome (IBS)

In medical literature, a description of IBS-type symptoms can be found as early as 1820¹. The symptoms included “occasional pain in the intestines and derangement of their powers of digestion, with flatulence, and a sense of suffocation”. Since that, these symptoms have been given names such as “spasmodic stricture of the colon”², “mucous colitis”³, “neurogenic mucous colitis”⁴, “colonic spasm”⁵, and “irritable colon syndrome”⁶. Nowadays, IBS is defined as “a functional bowel disorder in which abdominal pain or discomfort is associated with defecation or a change in bowel habit, and with features of disordered defecation”⁷. The term “functional” refers to absence of any organic disease, or structural or biochemical abnormality causing the symptoms. Abdominal pain or discomfort relieved by defecation indicates a possible colonic source, and appearance of pain associated with a change in bowel habits is indicative of a change in intestinal transit time, which may be a result of a change in intestinal motor or secretory function⁸.

Symptoms of IBS follow a chronic relapsing course. After one to ten years of follow-up, 50 to 70% of subjects still fulfil IBS criteria⁹⁻¹¹. The mean duration of GI symptoms among current IBS sufferers ranges between 5 and 13 years. A long history of GI symptoms and ongoing life stress are predictors of persisting symptoms⁸.

IBS is a benign disorder with no reported increased mortality in itself¹². IBS is, however, associated with an increased level of abdominal surgery^{13,14}, which may, together with unnecessary medical procedures, produce a source of elevated mortality rate not detected in studies so far¹⁵. Despite a benign nature, IBS reduces the quality of life (QoL) similarly to organic diseases, such as gastro-oesophageal reflux disease (GERD), asthma, or migraine¹⁶.

Pathogenesis

The pathogenesis of IBS is not fully understood. Abnormal motor function, visceral hypersensitivity, abnormal central nervous processing of visceral stimuli, gastrointestinal infections, subtle inflammation, psychosocial factors, abnormal gas handling, alterations in gut microflora, and genetic factors are possible agents involved in the pathogenesis of symptoms in IBS¹⁷.

The enteric nervous system (ENS) is a neural network coating the GI tract. Together with the sympathetic and parasympathetic system it forms the autonomic nervous system. The ENS integrates the contraction of smooth muscles, intestinal transport of water and electrolytes, intestinal secretion, and intramural blood flow. It contains sensory and motor neurons, interneurons and several neurotransmitters, such as serotonin (5-hydroxytryptamine; 5-HT), acetylcholine, opioids, noradrenaline, somatostatin,

cholecystokinin, substance P, and vasoactive intestinal polypeptide. It functions semi-autonomously, receiving input from the motor outflow of the sympathetic and parasympathetic systems and sending sensory information to the central nervous system (CNS)^{18,19}. According to a biopsychological conceptualisation, the brain-gut interaction between psychosocial and physiological factors influence GI symptom generation and outcome in terms of perceived symptom severity and medical care seeking²⁰ (Figure 1).




Figure 1. IBS conceptual model. IBS, irritable bowel syndrome; CNS, central nervous system; ENS, enteric nervous system.

Proximal colonic transit is accelerated in IBS with diarrhoea (IBS-D)²¹. In idiopathic constipation, both slowing of the whole colon and left colon transit is reported²². Also, altered small bowel transit with a correlation of GI symptoms²³ and abnormal response to stimuli, such as food and fatty acids, have been described²⁴.

IBS patients have a lower visceral pain threshold for both mechanical distension²⁵⁻²⁷ and electrical stimuli^{28,29} than controls, suggesting that primary afferent neurones (PAN) of the enteric nervous system are hypersensitive to non-noxious stimuli³⁰. Some studies indicate that they also have cutaneous hyperalgesia^{28,31,32}.

Functional magnetic resonance imaging (fMRI) and proton emission tomography (PET) studies show aberrant brain activation during noxious rectal distention and in the anticipation of rectal pain among IBS patients³³⁻³⁵, suggestive of abnormal central processing of the visceral stimuli. In an experimental fMRI study, amitriptyline reduced pain related cortical activation in IBS patients under stressful conditions³⁶. In a PET study,

cognitive therapy reduced limbic activity, GI symptoms, and anxiety³⁷, suggesting that cortical processing influences IBS symptoms. In addition, rectal stimulation at distension pressures below conscious perception show abnormal CNS activation in IBS patients, suggesting hypersensitivity of the neural circuitry irrespective of stimulus-related cognition³⁸.

Depression, anxiety, psychological distress, and stressful life events are more common among subjects with unexplained medical symptoms and increased use of healthcare services³⁹. Regardless of healthcare seeking status, an association between IBS and psychiatric distress has been established⁴⁰⁻⁴². In addition, over-representation of physical or sexual abuse has been reported in IBS patients⁴³⁻⁴⁵. Psychiatric disorders, especially anxiety, may precede the occurrence of GI symptoms and thus have a pathogenetic role in the development of IBS, with a link to affective spectrum disorders^{46,47}.

Infectious gastroenteritis precedes the onset of IBS in 6 to 17% of IBS patients⁴⁸. The relative risk of developing IBS after infection increases 10 to 12 times compared to uninfected controls^{49,50}. An increased number of mucosal lymphocytes, EC cells, higher levels of proinflammatory cytokines, and increased gut permeability are associated with post infective IBS⁵¹⁻⁵⁴. In addition, regardless of a history of gastroenteritis, neuronal degeneration in the jejunal myenteric plexus has been detected⁵⁵, as well as activation of the mucosal immune system, especially in IBS-D^{56,57}. Inflammatory changes in IBS and functional dyspepsia have even been reported in duodenal mucosa⁵⁸. A recent study found elevated levels, similar to those in active ulcerative colitis, of faecal human β -defensin-2 in IBS-D and mixed IBS (IBS-M) patients compared to healthy controls, supporting an activation of the mucosal innate defence system⁵⁹. IBS patients also have an increased frequency of activated T and B cells in their blood, consistent with a low grade inflammation^{60,61}. One study reported a correlation between the severity of abdominal pain and activated mast cells in close proximity to mucosal nerve endings, suggestive of a peripheral mechanism of pain⁶². Moreover, IBS-D patients have displayed enhanced proinflammatory cytokine release associated with GI symptoms and anxiety⁶³.

Abdominal bloating, reported by up to 96% of IBS patients, is often the most bothersome symptom⁶⁴. IBS patients have shown impaired gas transit, without signs of excess intra-abdominal gas⁸. Bloating alone is associated with visceral hypersensitivity, while patients with bloating and an increase in girth have normal sensory thresholds suggesting different pathogenetic mechanisms of the symptoms⁶⁵. Bloating alone is reported more in IBS-D, while bloating with an increment in abdominal girth in IBS with constipation (IBS-C)⁸.

Microbial genome analysis has revealed differences in the intestinal microbiota between IBS patients and healthy controls as well as between IBS subgroups, suggesting that intestinal bacteria also play a role in IBS development^{30,66}.

Genetic factors may influence development of IBS. Subjects who have a family member with GI symptoms are at a more than twofold risk of IBS-type symptoms, but having a spouse with these symptoms does not increase the risk⁶⁷. Genetics appear to have a role in functional bowel disorders, since a concordance of 33% between monozygotic twins and 13% between dizygotic twins has been reported, referring to a 57% share for genetics and 43% for environmental factors in a model by Morris-Yates *et al.*⁶⁸. Another

study concluded that social learning has an equal, or greater, influence than genetics on IBS development⁶⁹. A genetic component has also been reported for extra-intestinal symptoms associated with IBS⁷⁰. In addition, genetic polymorphism of serotonin transporter (SERT) may have a role in GI symptom pathogenesis^{71,72}.

Diagnostic criteria

Diagnostic tests are performed to rule out organic diseases that may produce similar symptoms to IBS. Negative exclusion diagnosis by means of endoscopic, radiographic, and laboratory investigations is, however, costly and inconvenient for the patient. In order to turn IBS diagnosis into a positive symptom based diagnosis, Manning *et al.* introduced the first symptom-based diagnostic criteria for IBS in 1978 (Table 1). The more of the six symptoms present, the more accurately patients with IBS were discriminated from those with organic disease. Three of the six symptoms were pain related. In their study, 31 of the 32 outpatients with IBS had abdominal pain, but no information of the duration or frequency was given⁷³. Two of the six symptoms were present in 94% of patients with IBS and 45% in patients with organic disease; the sensitivity was 94% and specificity 55%. Three or more criteria had a sensitivity of 84% and specificity of 76%⁷³. In epidemiological studies, the required number of Manning symptoms to fulfil IBS criteria has usually been two or three⁷⁴⁻⁷⁷. For population studies, a cut off of two Manning symptoms have been suggested, because of a lower prevalence of organic diseases in the general population than among outpatients⁷⁸.

In 1989, a multinational committee of clinical-investigators published first Rome criteria for IBS diagnosis⁷⁹, originally presented at the Thirteenth International Congress of Gastroenterology held in Rome in 1988. IBS was defined as “a functional gastrointestinal disorder attributed to the intestines and associated with symptoms of: (a) abdominal pain, and/or (b) disturbed defecation, and/or (c) bloatedness or distension”⁷⁹. Later, presence of abdominal pain was suggested as a requirement for the diagnosis, but the decision to do so was left to the investigator⁸⁰. Terms such as spastic colon or irritable colon were no longer recommended. These criteria were revised later and published as the Rome I criteria⁸¹ (Table 1). The main change from the 1989 criteria was the requirement of abdominal pain for IBS diagnosis. Rome I criteria consisted of abdominal pain related symptoms and non-pain related symptoms, later named the IBS supporting symptoms (Table 1).

The Rome I criteria were again revised, and the Rome II criteria published in 1999 by a Working Team⁸². The Rome II criteria were a committee consensus based on research results and expert opinion. The development process included reviewing and commenting by international experts. In the Rome II criteria, the second part, *i.e.* non-pain related symptoms of the Rome I criteria was deleted, due to poor clustering of these symptoms in factor analyses^{83,84}, their lower prevalence among males⁸⁵, and their partial inclusion in the first, pain-related part of Rome I criteria. In addition, “discomfort” was added to “pain”, and symptom duration was extended from three months to 12 months, with abdominal pain or discomfort present for at least 12 weeks (not necessarily consecutive weeks)

(Table 1). Rome I and Rome II criteria have been developed for both clinical practise and research purposes, such as epidemiological surveys, pathophysiology research, and therapeutic trials.

In 2006, the Rome II criteria were again revised employing a consensus approach, leading up to publication of the Rome III criteria by the Rome Working Team⁸⁶ (Table 1). The main change concerned the time perspective: the diagnostic criteria must be fulfilled for the last three months instead of 12 months. In addition, symptoms had to have begun at least 6 months before clinical presentation addressing the chronic nature of symptoms.

Despite of wide use of the Rome criteria in research, their accuracy has received little study. For Rome I criteria, the sensitivity for detecting IBS was 85% and specificity 71% in a study of 602 patients⁸⁷. In a retrospective study, Rome I criteria had a sensitivity of 65% and specificity of 100%, and positive predictive value of 100% in the absence of alarm signs⁸⁸.

IBS subtypes

IBS can be divided into three subtypes according to predominant bowel habit. In IBS with constipation (IBS-C), stools are usually hard or lumpy, and bowel movement frequency is less than three per week. Straining during a bowel movement is common. In IBS with diarrhoea (IBS-D), stools are usually loose, mushy, or watery and bowel movements take place more than three times a day. In addition, urgency (having to rush to bathroom) is a common phenomenon. Mixed IBS (IBS-M) applies to subjects expressing both constipation and diarrhoea variably, for at least 25% of the bowel movements.⁷ After a period of constipation, frequency of bowel movements typically increases leading to a period of diarrhoea. Usually these periods fluctuate rapidly within hours or a up to week^{89,90}. Unsubtyped IBS refers to those not meeting the definition for IBS-C, IBS-D, or IBS-M, while Alternating IBS is recommended to define those whose IBS subtype changes to another over a longer, more than one year, period of time⁸⁹.

Each IBS subtype (IBS-D, IBS-C, IBS-M) forms about one third of the patients fulfilling the Rome II IBS criteria⁸⁹. The proportion of the mixed subtype has varied between 23%⁹¹ and 63%⁹² in population based studies, however, and also IBS-D has been reported to be the commonest subtype⁹³. During a three month follow-up, approximately half of IBS patients shift from IBS-C or IBS-D to IBS-M, but shifting from IBS-D to IBS-C, or vice versa, is less common⁹⁴.

The Rome II subtyping of IBS is based on non-pain related IBS supporting symptoms, while Rome III subtyping is based solely on stool consistency according to the Bristol Stool Form, which has shown good correlation with whole-gut transit time⁹⁵⁻⁹⁸. The validity of IBS subtypes is uncertain. In a comparison study, IBS subtypes defined by Rome II and Rome III criteria showed poor agreement⁹⁹.

Table 1. Manning, Rome I, Rome II and Rome III criteria for IBS.

Manning criteria for IBS (Usually 2 or 3 symptoms required for criteria fulfilment)

- Looser stools at onset of pain
- More frequent bowel movements at onset of pain
- Pain eased after bowel movement
- Visible (abdominal) distension
- Passage of mucus
- Feeling of incomplete evacuation

Rome I criteria for IBS

At least 3 months continuous or recurrent symptoms of:

1. Abdominal pain or discomfort which is:
 - Relieved with defecation; and/or
 - Associated with a change in frequency of stool; and/or
 - Associated with a change in consistency of stoolAnd
2. Two or more of the following, on at least a quarter of occasions or days:
 - Altered stool frequency,
 - Altered stool form (lumpy/hard or loose/watery),
 - Altered stool passage (straining or urgency, feeling of incomplete evacuation)
 - Passage of mucus,
 - Bloating or feeling of abdominal distension.

Rome II criteria for IBS

At least 12 weeks or more, which need not be consecutive, in the preceding 12 months of abdominal discomfort or pain that has two out of three features:

1. Relieved with defecation; and/or
2. Onset associated with a change in frequency of stool; and/or
3. Onset associated with a change in form (appearance) of stool.

Rome III criteria for IBS*

Recurrent abdominal pain or discomfort at least three days per month in the last three months associated with two or more of the following:

1. Improvement with defecation
2. Onset associated with a change in frequency of stool
3. Onset associated with a change in form (appearance) of stool

*Criterion fulfilled for the last three months with symptom onset at least six months prior to diagnosis

Epidemiology

Prevalence

The incidence of IBS shows a substantial variation ranging between 2 to 70 per 1 000 patient years¹⁰⁰⁻¹⁰⁴. In Western countries, the prevalence of IBS ranges between 3 and 25% of the population^{92,103,105}. In non-Western countries, both lower prevalence estimates¹⁰⁶⁻¹⁰⁸, and similar rates to Western countries have been reported¹⁰⁹.

Table 2 shows IBS prevalence in several population studies according to varying diagnostic criteria and gender. In general, the Rome II criteria give a lower prevalence estimate than the Manning criteria. Some studies have included only two of the six Manning symptoms, while others have required three symptoms. The more symptoms required, the lower the prevalence estimate of IBS, which partly explains the variation in prevalence estimates.

Sociodemographic factors

In most studies, female gender is over-represented amongst IBS populations by a ratio of 1.5 to 2.0^{77,100,105,110-114}. Some studies applying the Rome II criteria, however, report a female to male ratio closer to one¹¹⁵⁻¹¹⁸. IBS-C is more common among females than males, while IBS-D is more common among males¹¹⁹⁻¹²². Females more often report bloating and extraintestinal comorbidity¹²³.

Pain related symptoms (pain eased after bowel movement, pain related to change in bowel habits) are equally common in both sexes⁸⁴. Symptoms not related to pain (diarrhoea, constipation, bloating, extraintestinal manifestations), however, are more common among female patients^{85,123}. Inclusion of non-pain related symptoms may partly account for female over-representation in IBS by Manning criteria. According to some reports, Manning criteria are less reliable in males^{124,125}.

The prevalence of IBS is highest between 20 to 40 years old¹²⁶, and usually decreases with age^{105,127}.

Some studies report a decrease in IBS prevalence with increasing income^{93,128}, suggesting that IBS is more prevalent amongst the lower social class. On the other hand, affluent childhood has also been associated with adult IBS¹²⁹.

Table 2. Prevalence of IBS by varying diagnostic criteria

Study	Population	Country	Method	Response		IBS criteria	IBS prevalence		
				rate (%)	n		All	Male	Female
Talley <i>et al.</i> 1991 ⁷⁸	Olmsted County; random sample	USA	Postal survey	82	835	≥ 3M	12.8	12.1	13.6
Heaton <i>et al.</i> 1992 ⁷⁷	Physicians' registry sample	UK	Postal survey	72	1 896	≥ 3M	9.5	5.0	13.0
Jones <i>et al.</i> 1992 ¹¹⁰	Physicians' registry sample	UK	Postal survey		1 620	≥ 2 M	21.6	18.7	24.3
Drossman <i>et al.</i> 1993 ¹⁰⁵	Random sample of households	USA	Postal survey	66	5 430	Rome 1989	9.4	7.7	14.5
Österberg <i>et al.</i> 2000 ⁴²	Random suburban sample	Sweden	Postal survey	58	2 707	Rome I	10.6	7.4	13.3
Boyce <i>et al.</i> 2000 ⁷⁵	Random sample of electoral roll	Australia	Postal survey	72	2 910	≥ 2 M	13.6	9.8	17.2
						Rome I	4.4	2.2	6.4
						Rome II	6.9	4.6	9.2
Talley <i>et al.</i> 2001 ¹³⁰	Birth cohort of young adults	New Zealand	Questionnaire	86	890	≥ 2 M	12.7	10.8	14.6
						Rome II	4.3	3.3	5.3
Mearin <i>et al.</i> 2001 ⁷⁶	Random population sample	Spain	Personal interview	76	1 932	≥ 3 M	10.3	5.6	14.8
						Rome I	12.1	7.2	16.8
						Rome II	3.3	1.9	4.6
Bommelaer <i>et al.</i> 2002 ¹³¹	Random population sample	France	Postal survey	NR	11 131	Rome I	4.0	2.5	5.3
Saito <i>et al.</i> 2003 ¹¹⁵	Olmsted County; random sample	USA	Postal survey	72	643	Rome I	6.8	5.6	8.0
						Rome II	4.7	4.9	4.5
Hungin <i>et al.</i> 2003 ⁹²	Random population sample	8 European countries	Telephone interview	NR	41 984	Manning	6.5	NR	NR
						Rome I	4.2	NR	NR
						Rome II	2.9	NR	NR
Dapoigny <i>et al.</i> 2004 ¹³²	Random population sample	France	2 phase postal survey	76 and 83	15 120	Rome II	4.7	3.7	5.7
Wilson <i>et al.</i> 2004 ¹³³	Physicians' registry sample	UK	2 phase postal survey	62 and 78	4 807	Rome II	10.5*	6.6*	14.0*
Andrews <i>et al.</i> 2005 ⁹³	Household sample	USA	Internet survey	82	25 986	Rome II	6.6	4.7	8.2

≥ 2 M, At least two Manning symptoms

≥ 3 M, At least three Manning symptoms

*Including those with a previous IBS diagnosis by physician

NR, Not reported

Comorbidity in IBS

About half of the IBS patients in primary care have at least one comorbid somatic symptom, and up to 94% of them have a comorbid psychiatric disorder¹³⁴. The most frequently reported symptoms include fibromyalgia, headache, back pain, chronic pelvic pain, temporomandibular joint pain, pollakisuria, dyspareunia, heart palpitation, depression, anxiety, chronic fatigue syndrome, and somatoformic disorders. Most studies assessing comorbidity in IBS have been carried out in clinical settings, *i.e.* among health care users. In a population based study, IBS non-consulters have also demonstrated higher rates of psychiatric and somatic comorbidities than population controls¹³⁵.

Similarities occur in the demographic distributions and psychological profiles among IBS and many comorbid conditions, suggesting a possible common underlying pathophysiologic mechanism or a common underlying disorder with different manifestations, but the evidence of such factors is still missing¹³⁴.

Psychiatric comorbidity

Depression and anxiety are common societal conditions. In a pan-European study, a 6-month prevalence rate for depressive disorders was 17%¹³⁶. In the US, the life-time prevalence of major depression was 17% in the general population, and 25% had suffered an anxiety disorder¹³⁷. In a Finnish population based interview study, the 12-month prevalence of a major depressive episode was 9.3%¹³⁸.

A comorbidity of depression, anxiety, and medical illness is also common. In the WHO Collaborative Study, primary health care patients with anxiety disorders were nine times more likely to develop depression than those with no other illness. Compared to patients with two or more chronic medical conditions, those with anxiety were six times more likely to develop depression. In addition, 39% of patients with depression also had an anxiety disorder, and 44% of those with an anxiety disorder also had depression.¹³⁹

Patients with depression often present with overlapping somatic symptoms typically including medically unexplained pain³⁹. Almost 30% of patients with depression also meet the criteria of IBS^{140,141}.

Patients with depression, or anxiety, use more healthcare services than those without. Amongst high utilizers of primary healthcare, 24% suffer from major depression, 33% have a lifetime history of depression, and 40% generalised anxiety disorder. In total, 83% of the high utilizers have psychiatric condition, including panic disorder, at some time in their lives.¹⁴² Amongst patients with somatic medical conditions, the presence of a psychiatric condition, such as depression or bipolar disease, doubles the individual healthcare expenditures¹⁴³.

Mental distress and psychiatric symptoms are over-represented in IBS⁴². Common psychiatric diagnoses among IBS sufferers include generalised anxiety disorder, panic

disorder, post-traumatic stress disorder, and somatoform disorders including hypochondriasis and somatisation disorder¹⁴⁴⁻¹⁴⁶.

Depression and anxiety are more prevalent amongst subjects with IBS than population controls or patients with somatic gastrointestinal conditions, such as inflammatory bowel disease¹⁴⁷. Depression in IBS patients appears to be more common with females than males¹⁴⁸. In primary care, patients with at least one GI symptom have a four to five fold prevalence of severe depression or anxiety compared to patients with no GI symptoms¹⁴⁹.

Psychiatric comorbidity may partly account for the higher level of healthcare use reported in IBS. Anxiety is particularly associated with healthcare seeking behaviour¹⁰⁸. On the other hand, a strong association between IBS and psychiatric disorders have been found independently of healthcare seeking status¹⁵⁰.

Somatic comorbidity

Gastrointestinal comorbidity

Almost half of the IBS patients also manifest other GI disorders, such as functional dyspepsia (FD), GERD, functional constipation, and anal incontinence¹⁵¹. The separation of IBS, FD, and reflux has been criticised because of the poor clustering of these disorders in factor analyses and a strong tendency of subjects in epidemiological studies to flux between IBS, FD, reflux, and unspecified GI symptoms. Approximately 50% of subjects with GI symptoms such as IBS, functional dyspepsia, or reflux, change their symptom profile during a one year follow-up⁹. A possible common underlying mechanism or unspecific responses to pathophysiological and psychological disturbances have been proposed to explain the variety of functional GI symptoms⁹.

Up to 87% of the subjects with IBS also have FD^{9,152}. Among IBS patients in a tertiary referral centre, IBS-C shows a higher rate of comorbidity with FD than other subtypes¹⁵³. Co-occurrence of IBS and FD in primary care has been reported to increase referrals to secondary care¹⁵⁴.

GERD is a common symptom occurring in about 20% of subjects in the general population. IBS shows a substantial co-occurrence with GERD, as approximately 40% of IBS healthcare seekers also have symptoms of GERD¹⁵⁵.

Anal incontinence following first vaginal delivery has more frequently been reported (64%) in women with IBS compared to 10% without¹⁵⁶, possibly related to rectal hypersensitivity and hypocompliance, especially in IBS-D¹⁵⁷.

IBS has also been associated with a higher rate of abdominal surgery, especially subjects with psychiatric comorbidity. For example, cholecystectomy rates for IBS patients are three times as high as those for matched controls¹³.

Extraintestinal comorbidity

Several nonpsychiatric extraintestinal symptoms are more prevalent in subjects with IBS than amongst non-IBS controls. In a US study, costs due to extraintestinal comorbidity covered more than 65% of costs incurred by IBS¹⁵⁸. The most studied comorbidities are fibromyalgia, chronic fatigue syndrome, and chronic pelvic pain, all of which occur in 33 to 50% of IBS patients¹⁵¹. Population prevalence of fibromyalgia is 2%¹⁵⁹, chronic fatigue syndrome is 0.4%¹⁶⁰, and chronic pelvic pain occurs in 14% of females¹⁵¹.

Chronic headache, migraine, temporomandibular joint disorder, and back pain are also significantly more common amongst IBS patients than healthy controls¹⁵¹. Besides chronic pelvic pain, other urogenital symptoms, such as dysuria, interstitial cystitis, dysmenorrhea, premenstrual syndrome, and disturbances in sexual function have also been reported more frequently than expected. Other symptoms include urinary stones, cardiac palpitations, and bronchial asthma or hyper-reactivity¹⁶¹.

Many common underlying physiological mechanisms, such as smooth muscle disorder, immune dysfunction, inflammatory condition, neuromuscular disease accompanied by visceral hypersensitivity, female hormonal alterations, behavioural response to pain, alterations in serotonergic system, and psychological factors are possible links between IBS and extraintestinal comorbidities. A theory of a global disease with symptoms covering multiple organs has not been proven to be the case. Multivariate analyses show comorbidities of IBS to be diseases of their own rather than a manifestation of a single disorder. IBS-sufferers with somatic comorbidities, however, possibly represent an IBS subgroup with psychological factors playing a common aetiological role.¹⁵¹

Treatment options

The objective of treatment in IBS is to alleviate symptoms and increase QoL. A good patient-physician relationship is important for satisfactory treatment. The strength of the physician-patient relationship is inversely proportionate to the number of physician visits; a positive interaction between the physician and the patient has been associated with a reduced use of healthcare services^{12,162}. Providing a diagnosis for the patient with an explanation of the benign nature of the symptoms will reduce fear of a malignant disease, help him or her to cope with the symptoms, and may even reduce the need for pharmacological treatment¹⁶³⁻¹⁶⁵. Factors worsening or triggering the symptoms, such as psychosocial stress or diet should be reviewed. In primary care, patients often attribute their symptoms to stress, but in secondary care are more likely to have psychiatric comorbidity, and consider stress unimportant for symptoms^{166,167}. Patients seem to expect more benefit from advice for diet, lifestyle, and exercise than from drugs^{168,169}.

A total of 60 to 70% of subjects with IBS associate their GI symptoms with food sensitivity^{170,171}. Dietary triggers reported to exacerbate IBS symptoms include caffeine, lactose, alcohol, fatty food, wheat, corn, citrus, food rich in carbohydrates, and hot spices¹⁷¹⁻¹⁷³. Especially caffeine and lactose may exacerbate symptoms among subjects

with IBS-D. A poor correlation, however, exists between reported lactose intolerance and a true lactose malabsorption^{174,175}. In addition, lactose restriction among subjects with IBS, reporting symptom exacerbation after lactose ingestion, does not necessarily improve GI symptoms^{176,177}. Excess ingestion of fructose has also been associated with exacerbation of symptoms and occasionally a restriction diet may prove helpful^{178,179}. No allergic IgE mediated reaction has been found to explain symptom exacerbation after meals^{170,180}. Positive findings in some studies exploring the efficacy of food elimination diets may partly be due to a placebo effect, and due to lack of evidence, such diets are not routinely advised¹⁸¹.

Increasing dietary fibre is generally recommended, though proof of efficacy of symptom alleviation is limited. Fibre products accelerate stool transit, and they are effective for treating constipation in IBS, but not pain or diarrhoea^{182,183}. Moreover, insoluble fibre, such as wheat bran can even worsen symptoms such as abdominal bloating and flatulence^{165,184-186}. For global symptom relief in IBS, soluble ispaghula husk seems to increase the rate of adequate relief and alleviate symptom severity, but insoluble fibre is no more effective than placebo¹⁸⁶⁻¹⁸⁸.

Probiotics are live microorganisms capable of exerting health benefits on the host. Possible mechanisms of action include modulation of the host immune system, stimulation of defensive systems, and competitive decrease in the number of pathogens. Health benefits depend probably on the bacterial strain¹⁸⁹. Bifidobacteria and probiotic combinations improve IBS symptoms, particularly flatulence and bloating, while lactobacilli alone does not seem efficient^{181,190}. Optimal dosage and combination of probiotics are unknown at the time.

Pharmacological treatment

The drugs available for IBS at the moment have limited evidence of efficacy. Less than one quarter of IBS patients demonstrate complete relief of IBS symptoms⁹². In addition, side effects of pharmacological therapies may have a considerable negative life impact¹⁹¹. The placebo effect is particularly strong, up to 70% in IBS drug studies¹⁹². Drug treatment is targeted to the dominant symptom or symptoms, usually diarrhoea, constipation, abdominal pain, or bloating.

Loperamide, a synthetic opioid, not transversing the blood-brain barrier, is effective in the treatment of diarrhoea by reducing stool frequency and improving stool consistency, but it is no more effective than placebo in reducing pain, bloating, and global symptoms of IBS¹⁹³⁻¹⁹⁶.

In addition to dietary fibre supplementation and commercial fibre preparations for constipation, osmotic laxatives, such as lactulose, or polyethylene glycol (PEG) are commonly used, although little studied in IBS. Amongst subjects with functional constipation, a low dose PEG may be more effective and produce less flatulence than lactulose¹⁹⁷. Amongst adolescents with IBS-C, treatment with a PEG laxative had a favourable effect on stool frequency, but not on pain intensity¹⁹⁸.

Smooth muscle relaxants and anticholinergic antispasmodics improve abdominal pain, but evidence of improving global IBS symptoms is not sufficient^{181,199,200}. Generally these are taken on as-needed basis. Side-effects include dry mouth, dizziness, blurred vision, and constipation. Also, peppermint oil may be superior to placebo in alleviating IBS symptoms, possibly by relaxing smooth muscle^{201,202}.

Tricyclic antidepressants (TCA) and selective serotonin reuptake inhibitors (SSRI) alleviate neuropathic pain^{203,204}. In IBS, TCAs have been used for abdominal pain. They have both anticholinergic and non-selective serotonin reuptake actions. The neuromodulatory and analgesic features, thought to act by inhibiting visceral hypersensitivity, usually appear at lower doses than those used in the treatment of depression^{36,205,206}. Anticholinergic side effects are similar to those with antispasmodics. Daily administration may bring out side effects even with low doses, however, the number needed to harm is 22²⁰³. According to a recent meta-analysis, the number needed to treat (NNT) is 4 (95% CI 3 to 8)²⁰⁷. A gradual introduction of TCA and a treatment duration of 6 to 12 months before dose tapering, has been suggested¹⁷³. Patients with IBS-D may benefit of TCAs more than other subtypes^{208,209}.

SSRIs have been widely used for treatment of anxiety, depression, and somatisation disorders²¹⁰. A few studies have assessed the role of SSRIs in IBS. Paroxetine improves QoL and overall well-being without a significant change in pain^{211,212}. Fluoxetine may alleviate abdominal pain in patients with rectal hypersensitivity²¹³, and increase the frequency of bowel movements besides relieving pain in IBS-C²¹⁴. According to a meta-analysis of five SSRI studies, NNT with SSRIs is 3.5 (95% CI 2 to 14)²⁰⁷.

Alocetron, a 5-HT₃ receptor antagonist, has high quality evidence from several studies on relieving global IBS symptoms in IBS-D with a NNT of 8. Due to serious side-effects, such as ischemic colitis and constipation-related surgery, however, it is not available in the EU market. The 5-HT₄ partial agonist tegaserod is more efficient than a placebo in relieving global IBS symptoms in female patients with IBS-C or IBS-M. The proportion of responders is between 5 and 19% higher than with placebo in several studies. Due to a ten-fold risk (frequency 0.11%) of cardiovascular events compared to controls, however, the drug has been withdrawn from market.¹⁸¹

Psychological therapies

Cognitive behavioural therapy^{208,215,216}, hypnotherapy^{217,218}, relaxation therapy (such as meditation)²¹⁹⁻²²¹, stress management²²², multi-component psychological therapy^{223,224}, and dynamic psychotherapy^{211,225} have been used to treat IBS. In a meta-analysis, the NNT of psychological studies was 4 (95% CI 3 to 5). Patients not responding to other IBS therapies could possibly benefit of psychological therapies, especially cognitive behavioural therapy. Most of the data is derived from small studies, however, and more large scale studies are needed to establish the role of psychological therapies in IBS.²⁰⁷

Healthcare use

IBS symptoms are one of the most common reasons for physician visits in primary care and referrals to gastroenterologists^{105,226}. In the US, functional GI disorders, such as constipation, dyspepsia, and IBS were among six leading GI physician diagnoses in outpatient visits in 2002²²⁷. Of those with IBS symptoms, 23 to 84% consult a primary care physician in a one year period^{127,131,228-232}. According to some studies, females are up to 3.3 times more prone to seek healthcare for GI symptoms^{105,233}, while others have found no gender difference in consultation behaviour^{77,229,231}.

Severity and the number of GI symptoms, presence of comorbid conditions, psychological factors (such as coping ability), depression or anxiety, and reduced QoL have been shown to predict healthcare seeking^{77,229,231,234-236}. Those with high levels of somatisation are more likely to be referred to secondary and tertiary centres¹⁶⁶ and they also exhibit more psychiatric symptoms and abnormal illness behaviour. In population surveys or patients with mild GI symptoms, however, the association of psychiatric symptoms and healthcare seeking seems to be weaker^{237,238}.

Besides consultations for GI symptoms, subjects with IBS make 2 to 3 times as many physician visits for non-GI reasons, and have increased rates of hospitalisations and surgical procedures^{105,239,240}.

In an Italian study, IBS-D was associated with an increased rate of physician visits²³². In another study from US, however, no difference was detected in healthcare seeking behaviour between IBS subgroups¹²⁷.

Abnormal illness behaviour

Besides multiple somatic complaints, inappropriate physician consultations for minor illnesses have been associated with IBS²⁴¹. Illness behaviour refers to how a person perceives abnormal bodily signals, *i.e.* symptoms of a disease and how he or she acts upon them. The decision of consulting a physician depends on knowledge on various symptoms and possible diseases behind them, the perceived intensity of symptoms, and individual health concerns²³⁵. In IBS, normal visceral stimuli and bodily sensations can be misinterpreted as signs of a disease. Symptom severity, worry of a serious illness, anxiety, and depression has been shown to discriminate IBS consulters and non-consulters²⁴². Also, beliefs about the efficacy of available treatments can influence consultation decisions. Finally, social support and individual coping skills can moderate healthcare seeking behaviour²³⁴.

Economic impact

The healthcare costs of IBS patients are more than 50% higher than controls^{243,244}. IBS mostly affects working age people leading to productivity loss due to sick leave. In a study comparing IBS costs between the UK, Germany, Switzerland and Portugal, the total

annual costs varied between 700 and 1 600€ per subject²⁴⁵. In a population based study, annual direct costs were 860€ per patient in France²⁴⁶. Costs incurred by Rome II criteria are reported to be higher than those by Rome I²⁴⁷.

The perspective of a cost evaluation may be that of an individual patient, a third-party payer (*i.e.* an insurance company or government), an individual hospital, or the society as a whole. In a societal perspective all expenditures are included regardless of the payer. Total expenditures consist of direct, indirect, and intangible costs. Direct costs are those related to use of healthcare services, medical investigations, and therapy. Indirect costs refer to productivity losses due to absenteeism from work and time spent at healthcare services. Intangible costs are incurred by pain, suffering, and reduced QoL, factors usually not evaluable in monetary terms.

Direct costs

In a review of US and UK studies about the economic impact of IBS, direct annual costs ranged widely between \$348 to \$8750 per subject²⁴⁸. Comparison between studies is difficult because of calculation differences for mean costs per subject. Some authors include only those with nonzero charges (*i.e.* healthcare users)²⁴³, others divide total costs by all participants even including non-healthcare users²⁴⁶. In addition, some only report GI related costs, while others include all costs irrespective of a particular diagnosis²⁴³. Different healthcare systems may also impede comparison. In a comparison between the UK and US, however, the use of healthcare facilities was largely similar despite differences in healthcare funding²⁴⁹.

In the US, direct costs of IBS have been estimated at approximately \$1.5 billion annually. In a comparison of 17 GI diseases in 1998, the highest direct costs were for GERD (\$9.3 billion), followed by gallbladder disease, colorectal cancer, peptic ulcer, diverticular disease, pancreatic disease, non-foodborne diseases, and chronic liver diseases before IBS. Pharmaceutical costs were markedly higher for GERD (\$5.9 billion) than IBS (\$0.08 billion), accounting for the presence of few effective drugs for IBS. In IBS, office visits covered the largest share (17%) of direct costs, but only 6% in GERD.²⁵⁰ Also in France, office visits formed the largest share of direct costs²⁴⁶.

Hospital inpatient costs for IBS have shown a wide variation between studies. Some studies report equal costs for IBS and control groups^{158,251}, some higher for IBS^{244,252}, others lower for IBS group²⁵³. The share of hospital inpatient costs have ranged between 7 and 70% of direct IBS costs²⁵⁴.

Diagnostic testing in IBS is performed to seek or rule out possible organic disease causing GI symptoms. In the US, almost one quarter of colonoscopies performed on patients under age 50, are for IBS symptoms²⁵⁵. A linear relationship has been detected between levels of somatisation and the amount of diagnostic testing for GI symptoms in IBS patients²⁵⁶. In a retrospective cohort analysis, colonoscopy or barium enema had been performed on 47% of patients with IBS²⁵⁷.

Indirect costs

Subjects with IBS incur indirect societal costs due to missed work days and impaired work performance because of GI symptoms. In a US bank employee study, IBS caused a reduction of 21% in work productivity, equivalent to working less than 4 days in a 5-day work week²⁵⁸. In another study, at least one third of IBS patients missed 1 working day monthly and 46% of them reported impairment at work because of GI symptoms²⁴⁹. IBS-employees with impaired work performance have more non-GI comorbidities, such as fibromyalgia, chronic fatigue syndrome, and urinary tract symptoms, than those with normal work performance despite IBS²⁵⁹.

3. AIMS OF THE STUDY

The main objectives of this thesis were:

1. To assess the prevalence of IBS
 - a. To evaluate IBS prevalence according to a variety of IBS definitions and diagnostic criteria
 - b. To assess the differences in symptom severity and health care use between subjects meeting differing IBS criteria
2. To evaluate the comorbidity in IBS and its impact on health care use
3. To investigate the connection of depressive and GI symptoms in the general population
4. To evaluate the societal costs of IBS

4. MATERIALS AND METHODS

Sample

The study was a two-phase postal survey (Figure 2). A random sample of non-institutionalised persons aged 18 to 64, was drawn from the National Population Registry. It comprised 2 490 men and 2 510 women with subjects' name, address, postal code, and age.

Only subjects, whose mother tongue was Finnish, were sampled. Those living in the Ahvenanmaa communal area were not included, as 93% of its inhabitants had Swedish as their mother tongue at the time of the study.

The sample size of 5 000 subjects was based on an assumption of 10% IBS prevalence in the population according to previous studies in Western countries, and a response rate of 70% (3 500 questionnaires for analysis). This allows subgroup analyses in the IBS group to detect a 5 to 10% unit frequency difference of a variable between an IBS group and non-IBS group with 5% precision and 90% confidence.

Questionnaires

Questionnaire I (Appendix A)

Questionnaire I, which was sent to a random sample of 5 000 subjects, contained 58 items: basic demographics (age, gender, marital status, education, and working hours), presence of a variety of somatic conditions diagnosed by a physician (lactose intolerance, celiac disease, atopic diseases, or bronchial asthma). In addition, a history of abdominal surgery, abdominal pain during childhood, history of upper or lower GI endoscopy, presence of any chronic medical condition, and regular medication were inquired (see Appendix A).

Items covering IBS diagnostic questions and those related to severity of abdominal symptoms were translated into Finnish from the Rome II Integrative Questionnaire, covering functional GI disorders from oesophagus to anorectum, designed primarily for epidemiological surveys¹⁹. According to the coding instructions, detection of the Rome II IBS criteria is based on six items including a question on the frequency of abdominal pain (q17), pain improvement with defecation (q20), questions on frequency of bowel movements (q21-22) and a change in stool form (q23-24) associated with abdominal pain. In addition, the Rome II Questionnaire includes nine items covering IBS supporting symptoms: abnormal frequency of bowel movements (q30-31), abnormal form of stools (q32-33), feeling of incomplete evacuation (q34), straining during a bowel movement (q35), urgency (q36), passage of mucus (q37), and abdominal bloating (q38). These symptoms are used to subclassify IBS into constipation-predominant and diarrhoea-predominant patterns¹⁹. They also allow IBS detection according to the

Manning and Rome I criteria (Table 1). In the Questionnaire I, Manning 2 criteria were defined as fulfilment of at least two of the six Manning symptoms and presence of abdominal pain more often than one day out of ten during the previous year. In Manning 3 criteria, at least three Manning symptoms were required. Rome I criteria were defined as a combination of abdominal pain more often than one day out of four, pain eased after bowel movement or associated with a change in bowel habits, and at least two of the five IBS supporting symptoms⁸¹ (Table 1). Dyspepsia was defined as presence of abdominal pain or discomfort (q17), localised in upper abdomen (q27-28), and not relieved with defecation (q20-24).

The response scale for all items to identify IBS and dyspepsia, as well as an item assessing the frequency of daily activities disturbed by pain, was a five grade frequency scale, which has later been proven to perform better than a binary scale²⁶⁰. Answering options were: 1) not at all or rarely; 2) occasionally; 3) often; 4) very often; 5) almost always. Time scales were defined as follows: *Occasionally*: more than one tenth of the time; *Often*: more than one quarter of the time; *Very Often*: more than one half of the time¹⁹. The severity of abdominal pain and discomfort during the previous year was assessed by the 4-grade Likert scale (mild, moderate, severe, and very severe).

The items translated into Finnish were then back-translated into English by an American-English translator. The back-translated items were compared with the original ones. Minor differences were noticed in the choice and order of words; such as “not at all or rarely” in the original was turned into “seldom or never” in the cross-translation. The cut-off points (choices such as “often”) in the frequency scale were identical in both questionnaires, however, suggesting that translation of the questionnaire into Finnish would not affect the number of IBS subjects identified.

Response options for questions covering the number of physician visits for GI and non-GI symptoms during the previous year were: 1) none; 2) one to two times; 3) three to five times; 4) six to ten times; 5) more than ten times. To approximate the average number of physician visits, they were later recoded as: 0, 1.5, 4, 8, and 11. Sick leave days during the previous year were assessed by five choices: 1) none; 2) one to three days; 3) four to six days; 4) seven to twenty days; 5) more than twenty days; 6) not employed. To approximate the mean number of sick leave days, these choices were recoded as: 0, 2, 5, 10, and 22.

Frequency of taking prescription medication and over-the-counter medication for GI symptoms was assessed by five options: 1) not at all; 2) less than once a month; 3) on average one to three times a month; 4) on average one to three times a week; 5) daily or almost daily.

Frequency of headaches and back pain were inquired by four options: 1) less than once a month; 2) on average one to three times a month; 3) on average one to three times a week; 4) daily or almost daily.

The final section of Questionnaire I consisted of a Finnish version of the 13-item Beck's Depression Inventory Short Form (BDI-SF)²⁶¹⁻²⁶⁴. The Finnish version (R-BDI-SF) includes 14 items: 13 for depressive symptoms and 1 for anxiety. It has been applied in population screening among Finnish adults and adolescents^{265,266} and has shown a good correlation with the original 21-item Beck Depression Inventory²⁶⁷. R-BDI-SF

score ranges from 0 to 36. Depressive symptoms are graded as mild (score 5-7), moderate (score 8-15), or severe (score 16 or more). Anxiety is scored as mild (score 1), moderate (score 2), or severe (score 3)²⁶⁴.

Pilot study

To assess the acceptability and item selection of the seven-page Questionnaire I, we conducted a pilot study in August 2001. The questionnaire with an explanatory letter was given to 30 consecutive outpatients (17 women, 13 men) visiting a gastroenterologist in the Department of Gastroenterology, Helsinki University Central Hospital. The patients displayed a variety of gastrointestinal disorders: functional GI disorders, inflammatory bowel diseases, and hepatic diseases. They were asked to fill in the questionnaire in the waiting room before or after the doctor's appointment and were encouraged to give comments on the questionnaire regarding the amount of items, contents of the questions, understanding of the questions, and suggestions for improving the questionnaire. Twenty-five patients returned the questionnaire. Completion of the questionnaire took approximately 12 minutes, which was regarded as a reasonable time by the patients. The questions were generally regarded as comprehensible and no changes were made to the questionnaire after the pilot study.

Questionnaire II (Appendix B)

Questionnaire II, which was sent to those fulfilling IBS criteria in Questionnaire I, included detailed questions on the duration of GI symptoms, the most bothersome symptoms, and location of abdominal pain. GI-related healthcare use was assessed by questions on the number of physician visits at various levels of healthcare organisation (primary care, occupational health care, secondary care, emergency room, private sector). In addition, the number of upper GI endoscopies, colonoscopies, barium enema, and abdominal ultrasound ever performed, were inquired, as well as consumption of medication for GI symptoms within nine pharmacological groups (antacids, histamine-2-receptor blockers, proton-pump inhibitors, pain killers [including non-steroidal anti-inflammatory drugs and paracetamol], antidiarrhoeals, laxatives, fibre products, antispasmodics, and prokinetics) during the previous three months. In addition, family history of inflammatory bowel disease, colorectal cancer, and celiac disease was inquired, (see Appendix B).

The number of physician visits during the previous year was inquired by six options: 1) none; 2) once; 3) twice; 4) three times; 5) four to six times; 6) more than six times. To approximate the average number of visits, "four to six times" was recoded as "five", and "more than six times" as "seven".

For use of medication, the answering options were: 1) not at all; 2) less than twice a month; 3) two to three times a month; 4) one to two times a week; 5) three to five times a

week; 6) daily or almost daily. To approximate the mean monthly use of medication these options were recoded as 0; 1; 2.5; 6; 16, and 24.

The final part of the Questionnaire II consisted of a validated Finnish version of the RAND 36-Item Health Survey 1.0, a generic health-related QoL measurement tool²⁶⁸.

For estimating the direct and indirect IBS costs, a societal perspective was obtained, *i.e.* all costs were taken into account regardless of the payer. Direct costs included those for physician visits, transportations, GI endoscopies, and medication. Indirect costs included expenses due to sick leave and time costs of physician visits and endoscopies.

The unit costs for physician visits, transportation, and time costs were derived from the National Research and Development Centre for Welfare and Health²⁶⁹. Costs of GI endoscopies were calculated as average prices between five university hospital districts of Finland. Medication costs were calculated based on a defined daily dose weighted with unit sales records within the ATC-4 group. Retail prices without VAT were derived from pharmacist price lists. Sick leave expenses, applying the human capital method²⁷⁰, were valued as average of lost earnings obtained from Statistics Finland²⁷¹.

Data collection

Questionnaire I was sent on the 18th of September 2001. The first reminder for non-respondents was sent on the 30th of October, and the final reminder on the 12th of December 2001.

Questionnaire II was sent to 591 subjects who met IBS criteria either by Manning ≥ 2 symptoms, or Rome II criteria in Questionnaire I. First mailing was on the 20th of February 2002. For non-responders, two reminders were sent 6 weeks apart.

All returned questionnaires were optically scanned and the data transformed into an Excel form. Statistical analyses were performed with NCSS-2000 software for Windows (NCSS Statistical Software, Kaysville, UT) and SPSS 13 software for Windows (SPSS, Inc, Chicago, IL, USA).

Control groups

In study I, four partially overlapping IBS groups were formed (Manning 2, Manning 3, Rome I, and Rome II). Non-IBS controls were subjects who did not meet the IBS criterion in question regardless of the possible reporting of GI symptoms and meeting another of the four IBS criteria.

In study II, IBS was defined as those fulfilling the Rome II IBS criteria. The control group included all subjects not meeting Rome II criteria.

In study III, frequencies of GI symptoms were compared between subjects reporting depressive symptoms and non-depressive controls. The control group was defined as subjects with an R-BDI-SF score of less than five.

In study IV, IBS related costs were compared between subjects fulfilling Rome II criteria and Manning 2 criteria. The control group consisted of those subjects fulfilling neither Rome II nor Manning 2 criteria.

Statistical methods

Prevalence figures for categorical variables were calculated with 95% confidence intervals (CI). Due to overlap of the four IBS criteria, 95% CIs were applied for comparison of frequencies of binary variables between the groups. Differences were considered statistically significant, if their 95% CIs did not overlap. Direct adjustment against Finnish population aged 18 to 64 was applied for age and gender adjustment of the prevalence figures by various IBS criteria and depressive symptoms.

In study I, concordance between different diagnostic criteria was calculated applying kappa-statistics (κ). A κ -value of 0.0 to 0.2 indicates slight agreement, 0.2 to 0.4 fair agreement, 0.4 to 0.6 moderate agreement, 0.6 to 0.8 substantial agreement, and a κ -value of 0.8 to 1.0 near perfect agreement²⁷².

Several variables with multiple categories, such as marital status, basic education, working hours, depression score, and anxiety, were dichotomized for comparison. The chi-square test was applied for categorical and T-test for dimensional variables. In study II, the Bonferroni correction for multiple testing was applied. For nonparametric comparison of medians, the Mann-Whitney's U-test was applied between two-group testing, and Kruskal Wallis test between three groups.

In study II, the distribution of comorbid conditions was compared between GI consulters and non-consulters among IBS group and controls. GI-consulters referred to those with at least one physician visit for GI symptoms during the previous year. A loglinear model was applied for comparison of categorical variables, and a two-way analysis of variance for continuous variables. To detect independent predictors of healthcare seeking for GI and non-GI reasons, a stepwise logistic regression analysis was performed with 26 variables: fixed variables (IBS, age, gender, basic education), working hours, employment status, marital status, GI conditions (dyspepsia, diarrhoea, constipation, severe abdominal pain, disturbance of daily activities because of GI symptoms, lactose intolerance, celiac disease, history of abdominal pain during childhood, history of abdominal surgery), allergic conditions (asthma, food allergy, allergic dermatitis, allergic rhinitis), other extra-GI conditions (headache, back pain, chronic medical condition), and psychiatric conditions (depression, anxiety and insomnia). In addition, interaction variables of each comorbid condition and IBS were generated and included in the regression model. For non-GI reasons, a majority of subjects in both the IBS and control groups had made at least one consultation during the previous year. Therefore, a consulter was defined as a subject with at least three non-GI visits.

In study III, independent factors associated with depressive symptoms were generated with a stepwise logistic regression analysis. Altogether 14 variables were included in the model: age, gender, educational level, marital status, employment status, working hours,

chronic medical condition, anxiety, IBS, dyspepsia, diarrhoea, constipation, frequent abdominal pain, and history of abdominal surgery.

In study IV, the non-parametric bootstrap technique²⁷³ was applied for comparison of the skewed cost data. The number of bootstrap replications was 10 000 and 95% confidence intervals were constructed by bootstrap percentile method where the lower and upper limits of the confidence interval are the 0.025 and 0.975 quantiles of the bootstrap distribution. The p-values for the differences between means were constructed by a randomisation test. The number of permutations in the tests was 10 000.

All p-values calculated were two-tailed. A p-value of less than 0.05 was considered statistically significant.

Ethics

The Ethics Committee of the Helsinki University Hospital evaluated the study protocol and stated that no ethical considerations by the committee are needed as the study is a postal survey.

5. RESULTS

For Questionnaire I, 5 000 letters were sent. Due to unknown addresses, 22 were returned. After two reminders, a total of 3 631 subjects participated in the study by returning the Questionnaire. A total of 20 questionnaires were returned unfilled. Questionnaire II was sent to 591 subjects fulfilling the Manning or Rome II IBS criteria in Questionnaire I. Of Phase II questionnaires, 8 went to unknown addresses. Altogether 499 filled and two unfilled questionnaires were returned (Figure 2).

The overall response rate for Questionnaire I was 73%. It was higher among females than males (79% vs. 67%). Mean respondent age was 42.1 years (95% CI 41.6-42.5), and non-respondents 38.7 years (95% CI 37.3-39.0). Of respondents, 55% were females (female to male ratio 1.2:1), whereas 39% of non-respondents were females. Among both male and female respondents, the youngest age group (18-24 years) was under-represented and oldest age group (55-64 years) over-represented (Table 2).

For Questionnaire II, the overall response rate was 86% (n=501). No gender difference emerged in any of the age groups. Mean respondent and non-respondent age was 42.2 (95% CI 41.1-43.4) and 37.4 (95% CI 35.0-40.0) years. The proportion of females among respondents and non-respondents was 64% and 66% (p = 0.781).

Table 2. Response rate (%)

Age (years)	Questionnaire I				Questionnaire II			
	Male n=1 639	Female n=1 991	Total n=3 631	p*	Male n=179	Female n=322	Total n=501	p*
18-24	53.5	73.5	63.3	<0.000	81.0	88.0	85.9	0.436
25-34	67.1	76.2	71.6	0.002	82.5	77.9	79.5	0.338
35-44	62.6	78.7	70.6	<0.000	76.0	84.8	81.7	0.196
45-54	69.5	79.5	74.6	<0.000	94.7	87.4	90.3	0.144
55-65	81.1	85.0	83.1	0.124	94.9	91.4	92.7	0.509
Total	67.0	78.9	72.9	<0.000	86.5	85.6	85.9	0.781

*Male vs. female

n = number of respondents




Figure 2. Flow chart of the study. Manning 2, at least two Manning symptoms; Manning 3, at least three Manning symptoms.

Prevalence of IBS (I)

At least one of the four IBS criteria was met by 600 subjects, which accounted for 16.5% of the respondents. The female to male ratio was 1.8:1. Manning 2 criteria represented the broadest IBS definition with a crude prevalence of 16.2% (95% CI 15.0-17.4%). After adjusting for age and gender, the prevalence was 15.9% (95% CI 14.7-17.1%). The most stringent criteria were Rome II, with a total of 184 respondents. Crude IBS prevalence was 5.1% (95% CI 4.4-5.8%). After adjusting for age and gender, the prevalence figure remained the same.

A significant gender difference appeared in both the Manning 2 and Manning 3 groups, with female to male ratio of 1.5:1 and 1.4:1, respectively. In Rome I and Rome II groups, no gender difference was evident (Table 3).

Table 3. Prevalence of IBS according to varying diagnostic criteria (% , 95% CI)

	Manning2 (n=587)	Manning3 (n=353)	Rome I (n=201)	Rome II (n=184)
Crude				
Total	16.2 (15.0-17.4)	9.7 (8.8-10.7)	5.5 (4.8-6.3)	5.1 (4.4-5.8)
Male	13.1 (11.4-14.8)	8.3 (7.0-9.7)	5.1 (4.0-6.2)	5.1 (4.0-6.2)
Female	19.2 (17.4-20.9)	11.2 (9.8-12.6)	6.1 (5.0-7.2)	5.3 (4.3-6.3)
Adjusted				
Total	15.9 (14.7-17.1)	9.6 (8.6-10.6)	5.6 (4.8-6.4)	5.1 (4.4-5.8)
Male	12.8 (11.1-14.4)	8.1 (6.7-9.5)	4.9 (3.8-6.0)	4.9 (3.8-6.0)
Female	19.1 (17.3-20.9)	11.2 (9.8-12.6)	6.2 (5.1-7.3)	5.2 (4.2-6.2)

In Manning 2 and 3 groups, the prevalence of IBS showed a declining tendency, although not statistically significant, from age 35 upwards (Table 4).

Table 4. Fulfillment of varying IBS criteria according to age (% , 95% CI).

	Manning2 (n=587)	Manning3 (n=353)	Rome I (n=201)	Rome II (n=184)
Age 18-24 (n=416)	17.8 (14.1-21.5)	10.3 (7.4-13.3)	6.0 (3.9-8.8)	5.3 (3.3-7.9)
Age 25-34 (n=683)	17.4 (14.6-20.3)	10.5 (8.2-12.8)	5.3 (3.7-7.2)	5.3 (3.7-7.2)
Age 35-44 (n=837)	16.7 (14.2-19.3)	9.7 (7.8-11.9)	5.6 (4.2-7.4)	4.9 (3.5-6.6)
Age 45-54 (n=941)	15.4 (13.1-17.7)	9.8 (8.0-11.9)	5.7 (4.3-7.4)	4.8 (3.5-6.3)
Age 55-65 (n=753)	14.5 (12.0-17.0)	8.6 (6.7-10.9)	5.6 (4.1-7.5)	5.3 (3.8-7.2)
p (χ^2 test)	0.445	0.776	0.991	0.982

The mean age for those fulfilling an IBS criterion did not differ from those not fulfilling. In addition, no difference was detected regarding marital status. In both Manning groups, a lower proportion of subjects had primary school or less as their basic education compared to those not fulfilling Manning criteria. No differences in basic education appeared in the Rome groups. In the Manning 3 group, a larger share had irregular working hours than those not fulfilling Manning 3 criteria.

Regardless of criteria applied, the diarrhoea-predominant subtype was the most common covering 41 to 57% of subjects fulfilling any IBS criterion. The proportion of constipation-predominant subtype ranged between 16 and 27%. In Rome II group, frequency of constipation did not differ from that of the total sample (Table 6.)

Overlap of IBS criteria (I)

A substantial overlap between various IBS criteria was detected. Manning 2 criteria was met by 97% of those fulfilling Rome II criteria. Only four subjects (2.2%) in the Rome II group exclusively met Rome II criteria (Table 5). Manning 2 criteria showed a substantial agreement with Rome I and Rome II criteria, with kappa-values of 0.45 and 0.42. Manning 3 criteria's agreement with Rome I and Rome II was slightly better, kappa-values were 0.53 and 0.52. A substantial agreement (kappa-value 0.78) was detected between Rome I and Rome II criteria.

Table 5. Overlap between diagnostic criteria. Number and percentage of subjects in column group also meeting upper row criteria.

	M2		M3		RI		RII		Only Column criteria	
	n	%	n	%	n	%	n	%	n	%
M2	587	100	0	0	195	33	179	30	183	31
M3	0	0	353	100	157	44	149	42	182	52
RI	195	96	157	77	204	100	153	75	8	4
RII	179	97	149	81	153	83	184	100	4	2.2

Table 6. Sociodemographic and symptom characteristics of total sample and each IBS criterion

	Total sample (n = 3 631)	Manning2 (n = 587)		Manning3 (n = 353)		Rome I (n = 204)		Rome II (n = 184)	
			p†		p†		p†		p†
Gender, n (%)			p1**		p1*		p1 = 0.108		p1 = 0.640
Male	1 639 (45.2)	209 (35.6)		132 (37.4)		81 (39.7)		80 (43.5)	
Female	1 991 (54.8)	378 (64.4)		221 (62.6)		123 (60.3)		104 (56.5)	
Mean age, years (SD)	42.3 (12.8)	41.4 (12.9)	0.066	41.5 (12.7)	0.239	42.5 (12.9)	0.844	42.2 (13.1)	0.888
Marital status, n (%)									
Married or common law marriage	2 411 (67.1)	398 (67.8)	0.686	248 (68.6)	0.536	138 (67.6)	0.860	120 (65.2)	0.580
Basic education, n (%)									
Primary school or less	863 (24.1)	112 (19.2)	*	67 (19.1)	0.020	48 (23.6)	0.867	40 (22.0)	0.486
High school graduate	1 182 (33.1)	192 (32.9)	0.921	119 (33.9)	0.722	55 (27.1)	0.063	50 (27.5)	0.100
Irregular working hours, n (%)	751 (21.2)	138 (24.0)	0.074	90 (26.1)	0.019	53 (26.6)	0.053	45 (25.1)	0.183
Diarrhoea, n (%)	521 (14.3)	242 (41.2)	**	168 (47.6)	**	110 (53.9)	**	104 (56.5)	**
Constipation, n (%)	548 (15.1)	133 (22.7)	**	66 (18.7)	0.046	54 (26.5)	**	30 (16.3)	0.63736
Dyspeptic symptoms, n (%)	288 (7.9)	172 (29.3)	**	107 (30.3)	**	98 (48.0)	**	83 (45.1)	**
Lactose intolerance, n (%)	344 (9.6)	129 (22.2)	**	70 (19.9)	**	53 (26.1)	**	44 (24.3)	**
Abdominal pain in childhood, n (%)	393 (11.1)	137 (23.8)	**	89 (25.8)	**	68 (34.0)	**	61 (33.3)	**
Allergic condition, n (%)	626 (18.1)	128 (23.0)	**	83 (24.6)	*	40 (20.8)	0.314	43 (24.9)	0.018
Asthma, n (%)	122 (3.5)	33 (5.8)	0.001	22 (6.4)	0.002	10 (5.0)	0.224	14 (7.8)	0.001
Headache, n (%)	370 (10.3)	123 (21.0)	**	82 (23.3)	**	54 (26.6)	**	50 (27.3)	**
Back pain, n (%)	568 (15.9)	173 (29.6)	**	113 (32.2)	**	80 (39.6)	**	64 (35.0)	**
Depressive symptoms, n (%)	602 (17.0)	175 (30.5)	**	120 (34.8)	**	84 (42.9)	**	70 (39.3)	**
Anxiety, n (%)	817 (22.8)	229 (39.4)	**	139 (39.6)	**	92 (45.3)	**	77 (42.1)	**
Insomnia, n (%)	630 (17.5)	155 (26.5)	**	98 (27.8)	**	66 (32.4)	**	59 (32.1)	**

†p-value vs. those not meeting the criterion

p1-value male vs. female

* < 0.01

** < 0.0001

GI symptom severity (I)

Presence of severe or very severe abdominal pain was reported by 44% (95% CI 36.6-51.1), 45% (95% CI 38.0-51.9), 30% (95% CI 25.2-34.8), and 27% (95% CI 23.7-30.9) of subjects in Rome II, Rome I, Manning 3, and Manning 2 groups. Daily activities were disturbed by abdominal pain often or very often by 64% (95% CI 57.2-71.1), 65% (95% CI 58.5-71.6), 33% (95% CI 27.9-37.7), and 29% (95% CI 25.3-32.6) of subjects in Rome II, Rome I, Manning 3, and Manning 2 groups (see Study I, Figure 2).

Prevalence of depressive symptoms (III)

According to R-BDI-SF, a total of 602 participants reported depressive symptoms corresponding to 17.0% (95% CI 15.7-18.2%) of the total sample, with no gender difference. Due to incompletely filled R-BDI-SF, depression score could not be rated for 82 subjects, who were excluded from the analysis. After age and gender adjustment, the prevalence of depressive symptoms was 16.7% (95% CI 15.4-17.9%). Independent predictors of depressive symptoms were anxiety, frequent abdominal pain (pain more often than 1 day of 4), diarrhoea, constipation, and a chronic medical condition. Higher basic education and marriage or cohabiting with a partner protected against depressive symptoms (see Study III, Table 2).

Of subjects with depressive symptoms, 11.6% (95% CI 9.1-14.2%) fulfilled Rome II IBS criteria, compared to only 3.7% (95% CI 3.0-4.4%) of the non-depressive control subjects. In addition, diarrhoea, constipation, and dyspepsia rates were higher amongst subjects with depressive symptoms.

Comorbidity in IBS (I,II)

Rates of dyspeptic symptoms, lactose intolerance, headache, back pain, depression, anxiety, and insomnia were higher for subjects meeting any IBS criterion than for controls. In addition, they had more often experienced abdominal pain during childhood. Presence of asthma or an allergic condition (food allergy, allergic dermatitis, or allergic rhinitis) was reported more often than controls in both Manning groups and the Rome II group, but no excess rates were reported in the Rome I group (Table 6).

Prevalence of depressive symptoms (depression score 5 or more according to R-BDI-SF) in the Rome II, Rome I, Manning 3, and Manning 2 groups was 39% (95% CI 32.1-46.5%), 43% (95% CI 35.9-49.8%), 35% (95% CI 29.8-39.8%), and 31% (95% CI 26.7-34.3%). Depressive symptoms were significantly more common in the Rome I group

than the Manning 2 group. In addition, anxiety and insomnia were more common amongst those meeting an IBS criterion than those not meeting it (Table 6).

Healthcare use (I,II,IV)

During the previous year, a higher proportion of subjects fulfilling any IBS criterion had made physician visits for both GI and non-GI symptoms than the total sample. Compared to subjects fulfilling either of the Manning criteria, a larger proportion of those fulfilling either of the Rome criteria reported visiting physicians for GI symptom (Figure 3). No difference emerged in non-GI visits between IBS groups.

The mean number of GI visits was 0.8 (95% CI 0.7-0.9) and 1.5 (95% CI 1.1-1.8) in the Manning 2 and in Rome II groups. In the control group (those fulfilling neither Rome II nor Manning 2 criteria), subjects reported on average 0.23 (95% CI 0.20-0.26) GI visits during the previous year. For non-GI visits, subjects in the Manning 2 group reported on average 3.0 visits (95% CI 2.7-3.2) and those in the Rome II group 3.2 visits (95% CI 2.8-3.6). Subjects in the control group reported on average 2.3 (95% CI 2.2-2.3) non-GI visits during the previous year (IV).




Figure 3. Visits to physicians for GI and non-GI reasons during the previous year.

Predictors of healthcare seeking (II, III)

In study II we compared subjects fulfilling Rome II IBS criteria to controls. In the control group, rates of lactose intolerance, diarrhoea, food allergy and headache were higher among GI consultants (those with at least one physician visit for GI reasons) than among non-consulters. However, in the Rome II group, rates of these conditions were lower among GI consultants (see Study II, Table 1). In addition, in the control group, rates of

depressive symptoms and anxiety were higher for GI consulters, but no difference emerged in the Rome II group between GI consulters and non-consulters.

When comparing the rates of dyspepsia, extra-GI somatic symptoms (headache, back pain, allergic conditions), or psychiatric symptoms (depression, anxiety, insomnia) between GI consulters and non-consulters in the Rome II group, only dyspepsia was more common among GI consulters (Figure 4).




Figure 4. Rates of comorbid symptoms in Rome II group between GI consulters and non-consulters.

In the logistic regression analysis, the strongest independent predictors of GI visits were GI symptoms, such as disturbing abdominal symptoms (OR 4.06, 95% CI 2.74-6.01), severe abdominal pain (OR 2.56, 95% CI 1.74-3.79), and dyspeptic symptoms (OR 2.30, 95% CI 1.62-3.27). Of the non-GI symptoms in the regression model, only insomnia (OR 1.57, 95% CI 1.21-2.06), headache (OR 1.43, 95% CI 1.04-1.97), and chronic illness (1.39, 95% CI 1.08-1.78) predicted healthcare seeking for GI symptoms. Co-occurrence of IBS and the variables in the regression model did not increase the probability of consulting a physician.

Presence of chronic illness (OR 3.16, 95% CI 2.62-3.82), back pain (OR 2.01, 95% CI 1.61-2.51), and depressive symptoms (OR 1.79, 95% CI 1.37-2.10) were the strongest predictors of non-GI visits. Of the GI symptoms in the regression model, only presence of dyspeptic symptoms (OR 1.64, 95% CI 1.20-2.25) and a history of abdominal pain in childhood (OR 1.39, 95% CI 1.07-1.81), but not IBS, predicted non-GI visits (see study II, Table III).

Depressive symptoms and health care seeking for GI reasons

In study III, GI related health care use was compared between subjects with depressive symptoms and non-depressive controls. At least one visit for GI reasons during the previous year was reported by 24% compared to 13% of controls ($p < 0.0001$). In addition, 18% of subjects with depressiveness had sick leave because of GI symptoms compared to 9% of controls ($p < 0.0001$).

Societal costs (IV)

In study IV, annual societal costs were compared between subjects fulfilling Manning 2 and Rome II criteria. Total GI related costs per subject were higher in the Rome II group (497€, 95% CI 382-621€) than in the Manning 2 group (295€, 95% CI 246-347€, $p < 0.001$). When allowing for population prevalences according to each criterion, the nationwide annual GI related costs were higher by Manning criteria than by Rome II criteria (154M€, 95% CI 128-181M€ vs. 82M€, 95% CI 63-103M€).

Direct costs covered two thirds of the total GI costs. The single most expensive cost component was the share of physician visits, accounting for approximately 60 % (59% to 63%) of the direct costs and 40 % (40% to 42%) of the total GI related costs in IBS. The distribution of GI costs was nearly identical in Manning 2 and Rome II groups (Figure 5).




Figure 5. Distribution of GI expenses in Rome II and Manning 2 groups.

Subjects fulfilling either IBS criteria more often used GI medication than controls. The share of medications was 13% to 14% of the total GI costs. Acid controlling medication formed 74% to 78% of the drug costs. Of medications for lower GI symptoms, bulking agents incurred the highest expense, which was, however, only 5% to 7% of the drug

expenses. Medication expenses per subject were higher in the Rome II group (63€, 95% CI 44-86€) than in the Manning 2 group (42€, 95% CI 33-52€, $p < 0.001$). In the Manning 2 group, total drug expenses per subject were slightly higher among females than males (see study IV, Table 3).

Annual direct costs of GI endoscopies per subject were 40€ (95% CI 30-50€) in the Manning 2 group and 58€ (95% CI 39-81€) in the Rome II group ($p < 0.001$), with no gender difference. At least one colonoscopy was in the history of 46% (95% CI 38.0-53.7) of the subjects in the Rome II group and 32% (95% CI 28.0-36.4) in the Manning 2 group. For those with at least one colonoscopy performed, the average number of procedures was 1.6 (95% CI 1.3-1.8) in the Rome II group and 1.5 (95% CI 1.4-1.7) in the Manning 2 group.

For non-GI visits, the direct individual costs were 242€ (95% CI 223-260€), 262€ (95% CI 228-296€), and 184€ (95% CI 177-190€) in the Manning 2, Rome II, and control groups. Nationally, the direct non-GI visit costs amounted to 126M€ by Manning, and 43M€ by Rome II criteria. Direct GI and non-GI expenditures altogether ranged between 98 M€ by Rome II and 230 M€ by Manning criteria.

The average number of sick leave days per subject for GI symptoms was 0.9 (95% CI 0.7-1.2) and 1.8 (95% CI 1.0-2.6) in the Manning 2 and Rome II groups, while controls reported on average 0.3 (95% CI 0.2-0.4) days off work. No gender difference was evident in any group. Individual costs of these days were higher in the Rome II group (147€, 95% CI 88-214€) than in the Manning 2 group (82€, 95% CI 60-107€, $p < 0.001$), (see study IV, Table 4).

GI related costs were also assessed by duration of GI symptoms by three categories: symptom duration of less than three years, three to ten years, and more than ten years. No difference was detected in direct or indirect GI related costs between subjects with different symptom duration. In addition, age group had no effect on total costs.

6. DISCUSSION

The prevalence of IBS shows substantial variation depending on the applied diagnostic criteria. Of the four IBS criteria studied, both Rome criteria are more stringent than the older Manning criteria and they identify an IBS population with more severe GI symptoms and more use of health care resources than those identified by the Manning criteria. IBS imposes a considerable burden for IBS sufferers and society. It is associated with high rates of psychiatric and somatic comorbidity and healthcare use for both GI and non-GI reasons. High population prevalence of IBS and increased use of healthcare resources incur high societal costs. Annually, individual GI related costs are almost twice as high by Rome II as Manning 2 criteria. Manning 2 criteria, however, identifies a three times larger IBS group owing to almost double the total societal GI related costs in Finland compared to Rome II criteria.

Methodological aspects

Our study was a large scale population based survey with high response rates lending support to the findings. A population based approach enables inclusion of subjects not using healthcare services. The Rome II Integrative Questionnaire is designed for epidemiological surveys, and R-BDI –SF, validated in Finnish²⁶⁷, has been widely used in surveys for detecting depressive symptoms.

As the Rome II Questionnaire was not available in Finnish, we translated the items used in our study. Translation may produce a source bias as the meaning of some expressions may differ between languages and cultures. Back-translation of the Questionnaire did not, however, reveal such biases suggesting that the translation process did not affect the number of identified IBS subjects.

For Questionnaire I, females were more prone to respond and consequently female to male ratio in the final data was 1.22:1, while at the time of the study, the ratio was 0.98:1 among Finnish inhabitants of age 18 to 64 years. Although the response rate was good overall, only 53.5% of males in the youngest age group responded possibly biasing results concerning them. The responders were on average 3.4 years older than non-responders which probably does not affect the results as the difference is considerably small and the total costs in the study were not higher for subjects in the oldest age groups.

No formal diagnoses of IBS or depression were confirmed by a physician as the study was a postal survey. Therefore, it is possible that some subjects fulfilling IBS criteria had an organic disease explaining GI symptoms. In the general population, however, functional bowel disorders are much more prevalent than organic diseases causing IBS-like symptoms⁹. Therefore, we assume that the IBS prevalence rates detected by our questionnaire are not significantly biased by organic diseases in the population. Although the BDI-SF is a self-reported depression symptom-rating scale and cannot substitute a diagnostic interview, it has shown a good correlation with moderate and severe depression in medical in-patients²⁷⁴.

In Questionnaire I, the average number of physician visits and work absenteeism days due to GI symptoms were estimated based on five-scale categorised items, which may bias the cost estimation of non-GI visits and the number of sick leave days. In Questionnaire II, the six-scale categorised items on GI visits offer more exact data as only three subjects in the Manning group and two in the Rome II group reported more than six GI visits.

Comparison of IBS criteria

In previous literature, the comparison of prevalence estimates and economic burden of IBS between individual countries and studies has been hampered by differences in IBS definition and criteria. In addition, composition of study population, research methodology, and sampling methods vary between studies. In our study, a substantial variation in the prevalence rates of IBS was seen depending on the applied diagnostic criteria, in line with two previous studies applying various IBS criteria on a population sample^{75,76}. Even when applying the same study methodology across eight European countries, however, a two-fold variation has been detected in IBS rates suggesting that true prevalence variations exist between individual countries, associated possibly with cultural and dietary habits⁹².

In this study, a three-fold difference in IBS prevalence rates was detected between the four IBS criteria. Manning 2 criteria were the most liberal and Rome II the most stringent. Two previous studies comparing Manning 2 and Rome II criteria obtained two to three-fold prevalence figures, in line with our results^{75,130}. In a systematic review targeting US population based studies applying only Rome definitions, IBS prevalence was in the same order as ours ranging between 5 to 10% with a pooled prevalence of 7% (95% CI 6-8%)²⁷⁵.

In our study, almost all subjects fulfilling Rome II criteria also fulfilled Manning 2 criteria. They formed a virtual subgroup of Manning 2 group with more severe abdominal pain, more disturbances of daily activities from symptoms, more use of healthcare services, and higher GI symptom related costs than those identified by just Manning criteria. This finding can partly be explained by the lesser requirement of pain in Manning criteria. Of the six Manning symptoms, only three are related to pain (looser stools at onset of pain, more frequent bowel movements at onset of pain, and pain eased after bowel movement). Abdominal pain disturbs daily activities and it predicts the use of the healthcare system which incurs economic expenses. According to the coding system in the Rome II Integrative Questionnaire, the required frequency of abdominal pain or discomfort to fulfil the criteria is “often”, which means more frequently than one day out of four. In the original Rome II criteria, the required frequency of abdominal pain is at least once a week for a period of 12 weeks during the previous 12 months. The reason for the higher requirement of pain in the Questionnaire is possibly to concentrate a more homogenous IBS group for research purposes and clinical trials, although this has not been commented on by the authors of the Rome II book¹⁹. In the widely used Rome II Modular Questionnaire, the required frequency of abdominal pain or discomfort is also “often”, meaning in the last three months “the symptoms were present during at least three

weeks (at least one day in each week)”¹⁹. This interpretation of “often” would generate an IBS prevalence of 9.3% in our population sample. In the Rome III Questionnaire, the required frequency of abdominal pain or discomfort is at least two to three days a month. In pathophysiology research and clinical trials, however, a frequency of at least two days a week during screening evaluation is recommended for subject eligibility²⁷⁶.

Rome II criteria have been criticised as too restrictive for research and practice⁷⁵. In addition, poor agreement between IBS diagnosis by general practitioners and Rome II criteria has been shown²⁷⁷. A clinician’s diagnosis is usually based on a global conception of a patient’s symptoms and clinical signs rather than on a standardised questionnaire. According to the Rome II definition, a high proportion of patients with IBS-type symptoms, but only occasional abdominal pain or discomfort, would fit into categories with no requirement of pain, such as functional constipation, functional diarrhoea, or unspecified functional bowel disorder²⁷⁸.

Depression and anxiety

The presence of depressive symptoms and anxiety is higher in the IBS population than among controls regardless of applied criteria. Half of the respondents fulfilling the Rome II criteria reported depressive symptoms or anxiety (II). Of those with at least one physician visit for GI symptoms, 43% had depressive symptoms, but these symptoms were not independent predictors of GI visits. Our results show that psychiatric comorbidity in IBS is not limited to healthcare seekers, but also exists among non-consulters (II). In two studies from Scandinavia, high frequencies of psychiatric distress and mood disorders were detected among both IBS consulters and non-consulters, supporting our findings^{42,135}.

Depressive symptoms are prevalent in the general population regardless of IBS. In our study, 17% of participants reported at least mild depressive symptoms. Prevalence of clinical depression in Finnish population studies have ranged between 5% to 9%^{138,279}. Differences between prevalence estimates are probably due to different study methodology as these studies used the University of Michigan Composite International Diagnostic Interview (UM-CIDI). Moreover, we used a cut-off level of 4/5 in R-BDI-SF, originally suggested for general population screening²⁶⁷, while a higher cut-off point of 7/8 has also been recommended to increase the specificity of R-BDI-SF for detection of depression among adolescents²⁸⁰. Thus, our results also include mild depressive symptoms.

Depressive symptoms were associated with a high rate of GI symptoms, and IBS was more than three times as common among subject with depressiveness. In addition, presence of depressive symptoms was associated with increased use of healthcare services and work absenteeism for GI reasons (III).

In former studies, high co-occurrence of psychiatric and GI symptoms has been observed among healthcare users with IBS²⁸¹⁻²⁸³. Our results suggest that overlap of depression and GI symptoms is also considerable in general population. Possible reasons for such overlap include common pathophysiological mechanisms regulating both mood

and brain-gut interactions^{284,285}, or a lower threshold of interpreting signals from GI tract as disturbing. No conclusions on causality, however, can be made from the present cross-sectional study.

Comorbidity

In the general population, several somatic and psychiatric symptoms are overrepresented among subjects fulfilling Rome II criteria of IBS.

Almost half of the subjects meeting Rome II IBS criteria reported symptoms consistent with dyspepsia, compared with only 6% of those without IBS. Earlier reports show a similar overlap with IBS and functional dyspepsia^{9,152,286}. This high co-occurrence is suggestive of a common pathophysiological basis, or only one disorder with different symptom manifestations.

Of those meeting the Rome II IBS criteria, 69% reported somatic symptoms, such as headache, back pain, allergy, or asthma compared to 35% among controls. In previous studies, chronic headache are reported by 23 to 35% and back pain by 28 to 81% of subjects with IBS^{287,288}, in line with our findings.

Some studies report an increased rate of asthma or bronchial hyper-responsiveness among subjects with IBS^{92,161,289}. In the present study, diagnosis of asthma was reported 2.4 times more frequently by subjects with IBS than by controls (II). IBS has also been associated with a positive methacholine test²⁹⁰, although not confirmed^{291,292}. Several theories of a common pathogenesis exist to explain the co-occurrence of asthma and IBS. A disorder affecting both bronchial and GI tract involved with smooth muscle, neuromuscular, or inflammatory system is possible, but needs further study. No significant difference was detected, however, in the frequency of food allergy in our study (II).

Healthcare seeking

IBS is associated with an elevated rate of physician visits for both GI and non-GI reasons. In our study, 48 to 49% of subjects meeting either Rome criteria had consulted a physician for abdominal symptoms during the previous year (I), in line with consultation rates in former studies^{229,231}. In a previous study from Finland, 75 to 81% of the population had made at least one physician visit for any reason during the previous year²⁹³. In our study, 80% of subjects in the control group had at least one non-GI visit, compared with an even higher share of 86 to 89% of those meeting any IBS criteria. The higher share of non-GI consultations in IBS is likely due to the high rates of non-GI comorbidity association.

Predictors of GI consultations

The main predictors for GI visits were disturbing abdominal symptoms, severe abdominal pain, dyspeptic symptoms, lactose intolerance, and IBS (II). Of the non-GI symptoms, only insomnia and headache, but not depression nor anxiety, were predictive for GI visits. This is in line with an Australian study by Talley *et al.*²²⁹, where pain severity, but not psychological comorbidity, was an independent predictor of GI visits.

In previous literature, the role of psychiatric symptoms as predictors of healthcare use is inconclusive. Anxiety, depression, somatisation, and lowered QoL are associated with increased healthcare use^{108,231,234,294}. In other population based studies, however, psychological factors did not explain healthcare seeking for abdominal reasons^{229,231}.

In our study, co-occurrence of IBS and other somatic or psychiatric symptoms did not elevate the likelihood of consulting for GI reasons. In fact, the combination of lactose intolerance and IBS lowered the probability of consultation, perhaps because of such individuals having already consulted earlier and attributing their GI symptoms as lactose ingestion not needing physician care. Overall, our results indicate that the predictors for consulting are not different for those with IBS compared to controls.

Predictors of non-GI consultations

Presence of IBS had no impact on consultation rates for non-GI visits. Young age and presence of a chronic medical condition were the strongest predictors for them. The need for physician consultations and statements for applying for a job, studying, or driver's license may explain young age as a predictor for non-GI visits. Of the GI disorders and symptoms listed (dyspepsia, IBS, diarrhoea, constipation, lactose intolerance, history of abdominal surgery, celiac disease, and abdominal pain), only the presence of dyspeptic symptoms predicted non-GI visits. One reason for dyspeptic symptoms predicting non-GI visits may be the patient's concern of possible serious disease, such as heart disease causing the upper abdominal symptoms²⁹⁵. It is also possible that dyspepsia is associated with other non-GI comorbidities, but not covered by our questionnaire, influencing healthcare use.

Societal costs

Due to high prevalence rates and frequent healthcare use, IBS generates a considerable economic burden. Our study indicates that 2 to 5% of the working age population annually consults a physician for their IBS symptoms. Those meeting Rome II criteria incur 1.7 times higher GI related costs per subject as those meeting the more lax Manning 2 criteria (497€ vs. 295€) (IV). Nationwide, however, expenditures are almost twice as high for Manning 2 as Rome II criteria (154M€ vs. 82M€), because of the considerable difference in the prevalence estimates by the criteria: 15.9% by Manning and 5.1% by Rome II

criteria. Overall, we obtained more modest individual GI costs compared to previous studies assessing IBS costs in Europe^{245,246}.

No gender difference was detected in GI related costs per subject, although some studies have found up to 3.3 times higher consultation rates for females^{105,116,233,296}. In the study by Drossman *et al.*¹⁰⁵, 44% of females and 35% of males with a functional GI disorder by Rome I criteria had visited a physician for GI symptoms during the previous year. In our study, 53% of females and 44% of males fulfilling Rome I criteria for IBS had made at least one physician visit for GI reasons ($p = 0.205$). In the database study by Shih *et al.*²³³, differences in healthcare use between genders were partly explained by gender differences in the population prevalence of IBS. Other studies from the UK, US, and Australia have, however, found no difference in consultation behaviour between genders^{77,229,231}. Likewise, two previous studies reported no gender difference in IBS costs^{243,246}. It is possible that cultural differences in healthcare use between genders in different countries exist, but this issue needs further study.

The principal cost component was the share of physician visits, covering 40 to 42% of the total annual costs. The average annual number of visits for GI symptoms ranged between 1.0 (Manning 2 criteria) and 1.8 (Rome II criteria). Higher number of annual physician visits for GI reasons have been reported by France (7.8)²⁴⁶, Spain (2.9)²⁴⁷, the UK (3.9)²⁴⁹, and the US (3.7)²⁴⁹. In the US Householder study¹⁰⁵, the number of GI visits was 1.64, closer to our findings.

Of the total GI expenditures, indirect costs covered 32 to 34% (IV). In previous studies, a wide range of 16 to 82% has been reported to represent the share of indirect costs in IBS. In a review, three out of five studies reported a proportion of more than 70% of the total costs. The main factor of indirect costs is sick leave cost. In our study, the mean number of sick leave days was 0.9 for those meeting Manning 2 criteria, considerably less than the 2.2 days in the French study²⁴⁶. Previous studies from the UK and US have reported even higher numbers of IBS-related sick leave days, ranging between 8.5 and 21.6 per year²⁴⁸. In our study, almost one third of subjects meeting Rome II criteria reported impaired performance at work weekly or more often indicating that they tend to go to work despite of GI symptoms rather than take sick leave. This may partly explain the lower than previously detected missed work days in our present study.

The duration of GI symptoms had no effect on total GI costs, or costs of endoscopies. IBS is a chronic disorder typically with fluctuating severity of symptoms. Repeating diagnostic procedures is not suggested, as the diagnosis is not likely to change over time^{81,297}. That the costs of endoscopies were not lower for those with several years of IBS symptoms may indicate that contrary to guideline recommendations, diagnostic procedures are repeated, or the first procedure is performed late in the course of GI symptoms. On the other hand, situations not covered by a cross sectional setting, such as a change in symptoms or appearance of new alarm symptoms, may also explain the finding.

In our present study, subjects meeting Manning 2 or Rome II criteria made 1.3 to 1.4 times more non-GI physician visits than controls. The direct costs of GI symptoms and non-GI visits amounted to 98M€ by Rome II and 230M€ by Manning 2 criteria, equivalent to a substantial share of 2 to 5% of the 4 562M€ expenditures on outpatient healthcare and medicines in Finland (IV). Besides direct and indirect costs, IBS incurs intangible costs

related to suffering from pain and reduced QoL, which cannot be properly measured in monetary terms. An integral approach to IBS by physicians, to also consider comorbid conditions, may produce a more favourable course in IBS patients and even reduce healthcare expenditures.

SUMMARY AND CONCLUSIONS

Our study investigated the prevalence of IBS according to varying diagnostic criteria, somatic and psychiatric comorbidity, healthcare resource use, and societal costs.

The prevalence of IBS shows a substantial variation depending on the criteria applied. Manning criteria with at least two of the six Manning symptoms represent the broadest interpretation of IBS giving a three-fold prevalence figure compared to Rome II, the strictest criteria. Subjects fulfilling Rome II criteria form a subgroup of those fulfilling Manning 2 criteria with more severe GI symptoms, more frequent health care resource use and higher societal costs per subject.

IBS, by Rome II criteria, is associated with a high level of somatic and psychiatric comorbidity among both healthcare consulters and non-consulters. Healthcare seeking for GI symptoms is predicted by symptom severity rather than psychiatric comorbidity. Presence of dyspeptic symptoms also predicts healthcare seeking for non-GI symptoms.

Depressive symptoms are prevalent in the general population and are associated with an increased level of GI symptoms, physician visits, and work absenteeism for GI symptoms.

The economic burden of IBS is considerable. Direct costs of GI symptoms and non-GI consultations correspond to a share of up to 5% of the outpatient healthcare and medicine expenditures in Finland.

Acknowledgements

This study, beginning in 2001, was carried out at the Division of Gastroenterology at Helsinki University Central Hospital.

I owe my deepest gratitude to Docent Martti Färkkilä who suggested this topic of study to me and encouraged me to enter the field of science. I also had the privilege to specialise in gastroenterology under his supervision. His profound experience and support during the several years of completing this thesis has been invaluable.

I am grateful to Professor Lars Agreus and Docent Rauli Leino, the official reviewers of this thesis, for their constructive comments and rapid review process.

I express my gratitude to my co-author Matti Siivola, MSc, who helped me through many statistical challenges. He also patiently taught me biostatistics during many sessions at the Department of Public Health and rapidly responded to my several bothersome e-mail consultations. I also thank Professor Seppo Sarna, Professor Pasi Korhonen, and Biostatistician Anna Wiksten for their invaluable statistical advice.

I owe my warmest thanks to Docent Martti Heikkinen and Juha Hämäläinen, MD, for their invaluable help in the field of psychiatric epidemiology and contribution as co-authors.

I am grateful to Niilo Färkkilä, MSc, for introducing me to the world of health economics and his enormous help in transforming our questionnaire data on healthcare use into monetary units. Moreover, his contribution as a co-author is greatly acknowledged.

I owe much to my colleges at Meilahti and Jorvi Hospitals for many fruitful discussions, practical advices, and friendly support to speed up the rather leisurely pace of my scientific work during these years. I especially thank Perttu Arkkila, Tarmo Koivisto, Jari Koskenpato, Henna Rautiainen, and Taina Sipponen for their time, constructive comments, hints, and ideas to various parts of this thesis.

I want to thank Graphic Designer Elina Nuortie for the cover of this book, and Software Specialist Anu Uusitalo for scanning the questionnaires and transforming the data onto an excel form.

This study was financially supported by the Finnish Foundation for Gastroenterological Research, the Mary and Georg C. Ehrnrooth Foundation, the Signe and Ane Gyllenberg Foundation, the Helsinki University EVO Foundation, the Finnish Cultural Foundation, and Novartis Finland OY. I am grateful for the trust of all those involved.

I owe my whole-hearted gratitude and appreciation to my parents, sister, and brother, together with my parents-in-laws for their continuous support and encouragement. Finally, my deepest gratefulness belongs to Anu and our children Petri and Taru for their love and patience during my years spent at the computer in the cellar of our house. All this time their support has carried me through the setbacks and strains of research work.

Espoo, February 2010

Markku Hillilä

References

1. Powell R. On certain painful afflictions of the intestinal canal. *Medical Transactions of the Royal College of Physicians* 1820;6:106-17.
2. Howship J. Practical remarks on the discrimination and successful treatment of spasmodic stricture in the colon. London: Burgess & Hill; 1830.
3. Osler W. The principles and practice of medicine: designed for the use of practitioners and students of medicine. New York: D Appleton; 1892.
4. Bockus HL, Bank J, Wilkinson SA. Neurogenic mucous colitis. *Am J Med Sci* 1928;176:813-29.
5. Ryle JA. Chronic spasmodic affections of the colon and the diseases which they simulate. *Lancet* 1928;ii:1115-9.
6. Chaudhary NA, Truelove SC. The irritable colon syndrome. *QJ Med* 1962;31:307-22.
7. Drossman D, Corazziari E, Delvaux M, Spiller RC, Talley NJ, Thompson WG, Whitehead WE, eds. Rome III: The Functional Gastrointestinal Disorders. Third edition. McLean, Virginia: Degnon Associates, 2006:490-509.
8. Spiller R, Aziz Q, Creed F, Emmanuel A, Houghton L, Hungin P, Jones R, Kumar D, Rubin G, Trudgill N, Whorwell P. Guidelines on the irritable bowel syndrome: mechanisms and practical management. *Gut* 2007;56:1770-98.
9. Agreus L, Svärdsudd K, Nyren O, Tibblin G. Irritable bowel syndrome and dyspepsia in the general population: overlap and lack of stability over time. *Gastroenterology* 1995;109:671-80.
10. Williams RE, Black CL, Kim HY, Andrews EB, Mangel AW, Buda JJ, Cook SF. Stability of irritable bowel syndrome using a Rome II-based classification. *Aliment.Pharmacol.Ther.* 2006;23:197-205.
11. Ford AC, Forman D, Bailey AG, Axon AT, Moayyedi P. Irritable bowel syndrome: a 10-yr natural history of symptoms and factors that influence consultation behavior. *Am.J.Gastroenterol.* 2008;103:1229,39; quiz 1240.
12. Owens DM, Nelson DK, Talley NJ. The irritable bowel syndrome: long-term prognosis and the physician-patient interaction. *Ann.Intern.Med.* 1995;122:107-12.
13. Longstreth GF, Yao JF. Irritable bowel syndrome and surgery: a multivariable analysis. *Gastroenterology* 2004;126:1665-73.
14. Kennedy TM, Jones RH. Epidemiology of cholecystectomy and irritable bowel syndrome in a UK population. *Br.J.Surg.* 2000;87:1658-63.
15. Talley NJ. Functional gastrointestinal disorders as a public health problem. *Neurogastroenterol.Motil.* 2008;20 Suppl 1:121-9.

16. Frank L, Kleinman L, Rentz A, Ciesla G, Kim JJ, Zacker C. Health-related quality of life associated with irritable bowel syndrome: comparison with other chronic diseases. *Clin.Ther.* 2002;24:675,89; discussion 674.
17. Barbara G, De Giorgio R, Stanghellini V, Cremon C, Salvioli B, Corinaldesi R. New pathophysiological mechanisms in irritable bowel syndrome. *Aliment.Pharmacol.Ther.* 2004;20 Suppl 2:1-9.
18. Wood JD. Enteric neuroimmunophysiology and pathophysiology. *Gastroenterology* 2004;127:635-57.
19. Drossman D, Corazziari E, Talley NJ, Thompson WG, Whitehead WE, eds. Rome II: The functional gastrointestinal disorders. 2nd ed. McLean, VA, USA: Degnon Associates; 2000.
20. Jones MP, Dilley JB, Drossman D, Crowell MD. Brain-gut connections in functional GI disorders: anatomic and physiologic relationships. *Neurogastroenterol.Motil.* 2006;18:91-103.
21. Vassallo M, Camilleri M, Phillips SF, Brown ML, Chapman NJ, Thomforde GM. Transit through the proximal colon influences stool weight in the irritable bowel syndrome. *Gastroenterology* 1992;102:102-8.
22. Stivland T, Camilleri M, Vassallo M, Proano M, Rath D, Brown M, et al. Scintigraphic measurement of regional gut transit in idiopathic constipation. *Gastroenterology* 1991;101:107-15.
23. Kellow JE, Phillips SF. Altered small bowel motility in irritable bowel syndrome is correlated with symptoms. *Gastroenterology* 1987;92:1885-93.
24. Drossman DA, Camilleri M, Mayer EA, Whitehead WE. AGA technical review on irritable bowel syndrome. *Gastroenterology* 2002;123:2108-31.
25. Whitehead WE, Engel BT, Schuster MM. Irritable bowel syndrome: physiological and psychological differences between diarrhea-predominant and constipation-predominant patients. *Dig.Dis.Sci.* 1980;25:404-13.
26. Kellow JE, Phillips SF, Miller LJ, Zinsmeister AR. Dysmotility of the small intestine in irritable bowel syndrome. *Gut* 1988;29:1236-43.
27. Whitehead WE, Palsson OS. Is rectal pain sensitivity a biological marker for irritable bowel syndrome: psychological influences on pain perception. *Gastroenterology* 1998;115:1263-71.
28. Drewes AM, Petersen P, Rossel P, Gao C, Hansen JB, Arendt-Nielsen L. Sensitivity and distensibility of the rectum and sigmoid colon in patients with irritable bowel syndrome. *Scand.J.Gastroenterol.* 2001;36:827-32.
29. Rossel P, Drewes AM, Petersen P, Nielsen J, Arendt-Nielsen L. Pain produced by electric stimulation of the rectum in patients with irritable bowel syndrome: further evidence of visceral hyperalgesia. *Scand.J.Gastroenterol.* 1999;34:1001-6.
30. Arebi N, Gurmany S, Bullas D, Hobson A, Staggs A, Kamm M. Review article: the psychoneuroimmunology of irritable bowel syndrome--an exploration of interactions between psychological, neurological and immunological observations. *Aliment.Pharmacol.Ther.*

2008;28:830-40.

31. Bouin M, Meunier P, Riberdy-Poitras M, Poitras P. Pain hypersensitivity in patients with functional gastrointestinal disorders: a gastrointestinal-specific defect or a general systemic condition? *Dig.Dis.Sci.* 2001;46:2542-8.
32. Verne GN, Robinson ME, Price DD. Hypersensitivity to visceral and cutaneous pain in the irritable bowel syndrome. *Pain* 2001;93:7-14.
33. Bonaz B. Visceral sensitivity perturbation integration in the brain-gut axis in functional digestive disorders. *J.Physiol.Pharmacol.* 2003;54 Suppl 4:27-42.
34. Silverman DH, Munakata JA, Ennes H, Mandelkern MA, Hoh CK, Mayer EA. Regional cerebral activity in normal and pathological perception of visceral pain. *Gastroenterology* 1997;112:64-72.
35. Bernstein CN, Frankenstein UN, Rawsthorne P, Pitz M, Summers R, McIntyre MC. Cortical mapping of visceral pain in patients with GI disorders using functional magnetic resonance imaging. *Am.J.Gastroenterol.* 2002;97:319-27.
36. Morgan V, Pickens D, Gautam S, Kessler R, Mertz H. Amitriptyline reduces rectal pain related activation of the anterior cingulate cortex in patients with irritable bowel syndrome. *Gut* 2005;54:601-7.
37. Lackner JM, Lou Coad M, Mertz HR, Wack DS, Katz LA, Krasner SS, Firth R, Mahl TC, Lockwood AH. Cognitive therapy for irritable bowel syndrome is associated with reduced limbic activity, GI symptoms, and anxiety. *Behav.Res.Ther.* 2006;44:621-38.
38. Lawal A, Kern M, Sidhu H, Hofmann C, Shaker R. Novel evidence for hypersensitivity of visceral sensory neural circuitry in irritable bowel syndrome patients. *Gastroenterology* 2006;130:26-33.
39. Katon W, Sullivan M, Walker E. Medical symptoms without identified pathology: relationship to psychiatric disorders, childhood and adult trauma, and personality traits. *Ann.Intern.Med.* 2001;134:917-25.
40. Drossman DA. Do psychosocial factors define symptom severity and patient status in irritable bowel syndrome? *Am.J.Med.* 1999;107:41S-50S.
41. Herschbach P, Henrich G, von Rad M. Psychological factors in functional gastrointestinal disorders: characteristics of the disorder or of the illness behavior? *Psychosom.Med.* 1999;61:148-53.
42. Österberg E, Blomquist L, Krakau I, Weinryb RM, Asberg M, Hultcrantz R. A population study on irritable bowel syndrome and mental health. *Scand.J.Gastroenterol.* 2000;35:264-8.
43. Drossman DA, Leserman J, Nachman G, Li ZM, Gluck H, Toomey TC, et al. Sexual and physical abuse in women with functional or organic gastrointestinal disorders. *Ann.Intern.Med.* 1990;113:828-33.
44. Leroi AM, Bernier C, Watier A, Hemond M, Goupil G, Black R, et al. Prevalence of sexual abuse among patients with functional disorders of the lower gastrointestinal tract.

Int.J.Colorectal Dis. 1995;10:200-6.

45. Talley NJ, Fett SL, Zinsmeister AR. Self-reported abuse and gastrointestinal disease in outpatients: association with irritable bowel-type symptoms. *Am.J.Gastroenterol.* 1995;90:366-71.
46. Solmaz M, Kavuk I, Sayar K. Psychological factors in the irritable bowel syndrome. *Eur.J.Med.Res.* 2003;8:549-56.
47. Sykes MA, Blanchard EB, Lackner J, Keefer L, Krasner S. Psychopathology in irritable bowel syndrome: support for a psychophysiological model. *J.Behav.Med.* 2003;26:361-72.
48. Longstreth GF, Hawkey CJ, Mayer EA, Jones RH, Naesdal J, Wilson IK, Peacock RA, Wiklund IK. Characteristics of patients with irritable bowel syndrome recruited from three sources: implications for clinical trials. *Aliment.Pharmacol.Ther.* 2001;15:959-64.
49. Parry SD, Stansfield R, Jelley D, Gregory W, Phillips E, Barton JR, Welfare MR. Does bacterial gastroenteritis predispose people to functional gastrointestinal disorders? A prospective, community-based, case-control study. *Am.J.Gastroenterol.* 2003;98:1970-5.
50. Rodriguez LA, Ruigomez A. Increased risk of irritable bowel syndrome after bacterial gastroenteritis: cohort study. *BMJ* 1999;318:565-6.
51. Gwee KA, Leong YL, Graham C, McKendrick MW, Collins SM, Walters SJ, Underwood JE, Read NW. The role of psychological and biological factors in postinfective gut dysfunction. *Gut* 1999;44:400-6.
52. Spiller RC, Jenkins D, Thornley JP, Hebden JM, Wright T, Skinner M, Neal KR. Increased rectal mucosal enteroendocrine cells, T lymphocytes, and increased gut permeability following acute *Campylobacter* enteritis and in post-dysenteric irritable bowel syndrome. *Gut* 2000;47:804-11.
53. Gwee KA, Collins SM, Read NW, Rajnakova A, Deng Y, Graham JC, McKendrick MW, Moolchala SM. Increased rectal mucosal expression of interleukin 1beta in recently acquired post-infectious irritable bowel syndrome. *Gut* 2003;52:523-6.
54. Dunlop SP, Jenkins D, Spiller RC. Distinctive clinical, psychological, and histological features of postinfective irritable bowel syndrome. *Am.J.Gastroenterol.* 2003;98:1578-83.
55. Törnblom H, Lindberg G, Nyberg B, Veress B. Full-thickness biopsy of the jejunum reveals inflammation and enteric neuropathy in irritable bowel syndrome. *Gastroenterology* 2002;123:1972-9.
56. Chadwick VS, Chen W, Shu D, Paulus B, Bethwaite P, Tie A, Wilson I. Activation of the mucosal immune system in irritable bowel syndrome. *Gastroenterology* 2002;122:1778-83.
57. Weston AP, Biddle WL, Bhatia PS, Miner PB, Jr. Terminal ileal mucosal mast cells in irritable bowel syndrome. *Dig.Dis.Sci.* 1993;38:1590-5.
58. Walker MM, Talley NJ, Prabhakar M, Pennaneac'h CJ, Aro P, Ronkainen J, Storskrubb T, Harmsen WS, Zinsmeister AR, Agreus L. Duodenal mastocytosis, eosinophilia and intraepithelial lymphocytosis as possible disease markers in the irritable bowel syndrome and

functional dyspepsia. *Aliment.Pharmacol.Ther.* 2009;29:765-73.

59. Langhorst J, Junge A, Rueffer A, Wehkamp J, Foell D, Michalsen A, Musial F, Dobos GJ. Elevated human beta-defensin-2 levels indicate an activation of the innate immune system in patients with irritable bowel syndrome. *Am.J.Gastroenterol.* 2009;104:404-10.
60. Öhman L, Isaksson S, Lindmark AC, Posserud I, Stotzer PO, Strid H, Sjövall H, Simren M. T-cell activation in patients with irritable bowel syndrome. *Am.J.Gastroenterol.* 2009;104:1205-12.
61. Öhman L, Lindmark AC, Isaksson S, Posserud I, Strid H, Sjövall H, Simren M. B-cell activation in patients with irritable bowel syndrome (IBS). *Neurogastroenterol.Motil.* 2009;21:644-50.
62. Barbara G, Stanghellini V, De Giorgio R, Cremon C, Cottrell GS, Santini D, Pasquinelli G, Morselli-Labate AM, Grady EF, Bunnett NW, Collins SM, Corinaldesi R. Activated mast cells in proximity to colonic nerves correlate with abdominal pain in irritable bowel syndrome. *Gastroenterology* 2004;126:693-702.
63. Liebrechts T, Adam B, Bredack C, Roth A, Heinzel S, Lester S, Downie-Doyle S, Smith E, Drew P, Talley NJ, Holtmann G. Immune activation in patients with irritable bowel syndrome. *Gastroenterology* 2007;132:913-20.
64. Houghton LA, Whorwell PJ. Towards a better understanding of abdominal bloating and distension in functional gastrointestinal disorders. *Neurogastroenterol.Motil.* 2005;17:500-11.
65. Agrawal A, Houghton LA, Lea R, Morris J, Reilly B, Whorwell PJ. Bloating and distention in irritable bowel syndrome: the role of visceral sensation. *Gastroenterology* 2008;134:1882-9.
66. Kassinen A, Krogius-Kurikka L, Mäkituokko H, Rinttilä T, Paulin L, Corander J, Malinen E, Apajalahti J, Palva A. The fecal microbiota of irritable bowel syndrome patients differs significantly from that of healthy subjects. *Gastroenterology* 2007;133:24-33.
67. Locke GR,3rd, Zinsmeister AR, Talley NJ, Fett SL, Melton LJ,3rd. Familial association in adults with functional gastrointestinal disorders. *Mayo Clin.Proc.* 2000;75:907-12.
68. Morris-Yates A, Talley NJ, Boyce PM, Nandurkar S, Andrews G. Evidence of a genetic contribution to functional bowel disorder. *Am.J.Gastroenterol.* 1998;93:1311-7.
69. Levy RL, Jones KR, Whitehead WE, Feld SI, Talley NJ, Corey LA. Irritable bowel syndrome in twins: heredity and social learning both contribute to etiology. *Gastroenterology* 2001;121:799-804.
70. Svedberg P, Johansson S, Wallander MA, Hamelin B, Pedersen NL. Extra-intestinal manifestations associated with irritable bowel syndrome: a twin study. *Aliment.Pharmacol.Ther.* 2002;16:975-83.
71. Camilleri M. Is there a SERT-ain association with IBS? *Gut* 2004;53:1396-9.
72. Yeo A, Boyd P, Lumsden S, Saunders T, Handley A, Stubbins M, Knaggs A, Asquith S, Taylor I, Bahari B, Crocker N, Rallan R, Varsani S, Montgomery D, Alpers DH, Dukes GE, Purvis I, Hicks GA. Association between a functional polymorphism in the serotonin

transporter gene and diarrhoea predominant irritable bowel syndrome in women. *Gut* 2004;53:1452-8.

73. Manning AP, Thompson WG, Heaton KW, Morris AF. Towards positive diagnosis of the irritable bowel. *Br.Med.J.* 1978;2:653-4.
74. Kay L, Jorgensen T, Lanng C. Irritable bowel syndrome: which definitions are consistent? *J.Intern.Med.* 1998;244:489-94.
75. Boyce PM, Koloski NA, Talley NJ. Irritable bowel syndrome according to varying diagnostic criteria: are the new Rome II criteria unnecessarily restrictive for research and practice? *Am.J.Gastroenterol.* 2000;95:3176-83.
76. Mearin F, Badia X, Balboa A, Baro E, Caldwell E, Cucala M, Diaz-Rubio M, Fueyo A, Ponce J, Roset M, Talley NJ. Irritable bowel syndrome prevalence varies enormously depending on the employed diagnostic criteria: comparison of Rome II versus previous criteria in a general population. *Scand.J.Gastroenterol.* 2001;36:1155-61.
77. Heaton KW, O'Donnell LJ, Braddon FE, Mountford RA, Hughes AO, Cripps PJ. Symptoms of irritable bowel syndrome in a British urban community: consulters and nonconsulters. *Gastroenterology* 1992;102:1962-7.
78. Talley NJ, Zinsmeister AR, Van Dyke C, Melton LJ,3rd. Epidemiology of colonic symptoms and the irritable bowel syndrome. *Gastroenterology* 1991;101:927-34.
79. Thompson WG, Dotevall G, Drossman DA, Heaton W, Kruis W. Irritable bowel syndrome: Guidelines for the diagnosis. *Gastroenterol Int* 1989;2:92-5.
80. Drossman DA, Thompson WG, Talley NJ, Funch-Jensen P, Janssens J, Whitehead WE. Identification of Sub-groups of Functional Gastrointestinal Disorders. *Gastroenterol Int* 1990;3:159-72.
81. Thompson WG, Creed F, Drossman DA, Heaton KW, Mazzacca G. Functional Bowel Disease and Functional Abdominal Pain. *Gastroenterol Int* 1992;6:75-91.
82. Thompson WG, Longstreth GF, Drossman DA, Heaton KW, Irvine EJ, Muller-Lissner SA. Functional bowel disorders and functional abdominal pain. *Gut* 1999;45:II43-7.
83. Whitehead WE, Crowell MD, Bosmajian L, Zonderman A, Costa PT,Jr, Benjamin C, Robinson JC, Heller BR, Schuster MM. Existence of irritable bowel syndrome supported by factor analysis of symptoms in two community samples. *Gastroenterology* 1990;98:336-40.
84. Taub E, Cuevas JL, Cook EW,3rd, Crowell M, Whitehead WE. Irritable bowel syndrome defined by factor analysis. Gender and race comparisons. *Dig.Dis.Sci.* 1995;40:2647-55.
85. Thompson WG. Gender differences in irritable bowel symptoms. *Eur.J.Gastroenterol.Hepatol.* 1997;9:299-302.
86. Longstreth GF, Thompson WG, Chey WD, Houghton LA, Mearin F, Spiller RC. Functional bowel disorders. *Gastroenterology* 2006;130:1480-91.

87. Tibble JA, Sigthorsson G, Foster R, Forgacs I, Bjarnason I. Use of surrogate markers of inflammation and Rome criteria to distinguish organic from nonorganic intestinal disease. *Gastroenterology* 2002;123:450-60.
88. Vanner SJ, Depew WT, Paterson WG, DaCosta LR, Groll AG, Simon JB, Djurfeldt M. Predictive value of the Rome criteria for diagnosing the irritable bowel syndrome. *Am.J.Gastroenterol.* 1999;94:2912-7.
89. Drossman DA, Morris CB, Hu Y, Toner BB, Diamant N, Leserman J, Shetzline M, Dalton C, Bangdiwala SI. A prospective assessment of bowel habit in irritable bowel syndrome in women: defining an alternator. *Gastroenterology* 2005;128:580-9.
90. Guilera M, Balboa A, Mearin F. Bowel habit subtypes and temporal patterns in irritable bowel syndrome: systematic review. *Am.J.Gastroenterol.* 2005;100:1174-84.
91. Mearin F, Balboa A, Badia X, Baro E, Caldwell E, Cucala M, Diaz-Rubio M, Fueyo A, Ponce J, Roset M, Talley NJ. Irritable bowel syndrome subtypes according to bowel habit: revisiting the alternating subtype. *Eur.J.Gastroenterol.Hepatol.* 2003;15:165-72.
92. Hungin AP, Whorwell PJ, Tack J, Mearin F. The prevalence, patterns and impact of irritable bowel syndrome: an international survey of 40,000 subjects. *Aliment.Pharmacol.Ther.* 2003;17:643-50.
93. Andrews EB, Eaton SC, Hollis KA, Hopkins JS, Ameen V, Hamm LR, Cook SF, Tennis P, Mangel AW. Prevalence and demographics of irritable bowel syndrome: results from a large web-based survey. *Aliment.Pharmacol.Ther.* 2005;22:935-42.
94. Mearin F, Baro E, Roset M, Badia X, Zarate N, Perez I. Clinical patterns over time in irritable bowel syndrome: symptom instability and severity variability. *Am.J.Gastroenterol.* 2004;99:113-21.
95. Heaton KW, O'Donnell LJ. An office guide to whole-gut transit time. Patients' recollection of their stool form. *J.Clin.Gastroenterol.* 1994;19:28-30.
96. Probert CS, Emmett PM, Heaton KW. Some determinants of whole-gut transit time: a population-based study. *QJM* 1995;88:311-5.
97. Degen LP, Phillips SF. How well does stool form reflect colonic transit? *Gut* 1996;39:109-13.
98. Lewis SJ, Heaton KW. Stool form scale as a useful guide to intestinal transit time. *Scand.J.Gastroenterol.* 1997;32:920-4.
99. Ersryd A, Posserud I, Abrahamsson H, Simren M. Subtyping the irritable bowel syndrome by predominant bowel habit: Rome II versus Rome III. *Aliment.Pharmacol.Ther.* 2007;26:953-61.
100. Kay L, Jorgensen T, Jensen KH. The epidemiology of irritable bowel syndrome in a random population: prevalence, incidence, natural history and risk factors. *J.Intern.Med.* 1994;236:23-30.
101. Locke GR,3rd, Yawn BP, Wollan PC, Melton LJ,3rd, Lydick E, Talley NJ. Incidence of a clinical diagnosis of the irritable bowel syndrome in a United States population.

- Aliment.Pharmacol.Ther. 2004;19:1025-31.
102. Agreus L, Svärdsudd K, Talley NJ, Jones MP, Tibblin G. Natural history of gastroesophageal reflux disease and functional abdominal disorders: a population-based study. *Am.J.Gastroenterol.* 2001;96:2905-14.
 103. Saito YA, Schoenfeld P, Locke GR,3rd. The epidemiology of irritable bowel syndrome in North America: a systematic review. *Am.J.Gastroenterol.* 2002;97:1910-5.
 104. Huerta I, Valdovinos MA, Schmulson M. Irritable bowel syndrome in Mexico. *Dig.Dis.* 2001;19:251-7.
 105. Drossman DA, Li Z, Andruzzi E, Temple RD, Talley NJ, Thompson WG, Whitehead WE, Janssens J, Funch-Jensen P, Corazziari E. U.S. householder survey of functional gastrointestinal disorders. Prevalence, sociodemography, and health impact. *Dig.Dis.Sci.* 1993;38:1569-80.
 106. Massarrat S, Saberi-Firoozi M, Soleimani A, Himmelmann GW, Hitzges M, Keshavarz H. Peptic ulcer disease, irritable bowel syndrome and constipation in two populations in Iran. *Eur.J.Gastroenterol.Hepatol.* 1995;7:427-33.
 107. Ho KY, Kang JY, Seow A. Prevalence of gastrointestinal symptoms in a multiracial Asian population, with particular reference to reflux-type symptoms. *Am.J.Gastroenterol.* 1998;93:1816-22.
 108. Hu WH, Wong WM, Lam CL, Lam KF, Hui WM, Lai KC, Xia HX, Lam SK, Wong BC. Anxiety but not depression determines health care-seeking behaviour in Chinese patients with dyspepsia and irritable bowel syndrome: a population-based study. *Aliment.Pharmacol.Ther.* 2002;16:2081-8.
 109. Masud MA, Hasan M, Khan AK. Irritable bowel syndrome in a rural community in Bangladesh: prevalence, symptoms pattern, and health care seeking behavior. *Am.J.Gastroenterol.* 2001;96:1547-52.
 110. Jones R, Lydeard S. Irritable bowel syndrome in the general population. *BMJ* 1992;304:87-90.
 111. Zuckerman MJ, Guerra LG, Drossman DA, Foland JA, Gregory GG. Comparison of bowel patterns in Hispanics and non-Hispanic whites. *Dig.Dis.Sci.* 1995;40:1763-9.
 112. Thompson WG, Irvine EJ, Pare P, Ferrazzi S, Rance L. Functional gastrointestinal disorders in Canada: first population-based survey using Rome II criteria with suggestions for improving the questionnaire. *Dig.Dis.Sci.* 2002;47:225-35.
 113. Wei X, Chen M, Wang J. The epidemiology of irritable bowel syndrome and functional constipation of Guangzhou residents. *Zhonghua Nei Ke Za Zhi* 2001;40:517-20.
 114. Rajendra S, Alahuddin S. Prevalence of irritable bowel syndrome in a multi-ethnic Asian population. *Aliment.Pharmacol.Ther.* 2004;19:704-6.
 115. Saito YA, Talley NJ, J Melton L, Fett S, Zinsmeister AR, Locke GR. The effect of new diagnostic criteria for irritable bowel syndrome on community prevalence estimates.

- Neurogastroenterol.Motil. 2003;15:687-94.
116. Lau EM, Chan FK, Ziea ET, Chan CS, Wu JC, Sung JJ. Epidemiology of irritable bowel syndrome in Chinese. *Dig.Dis.Sci.* 2002;47:2621-4.
 117. Lu CL, Chen CY, Lang HC, Luo JC, Wang SS, Chang FY, Lee SD. Current patterns of irritable bowel syndrome in Taiwan: the Rome II questionnaire on a Chinese population. *Aliment.Pharmacol.Ther.* 2003;18:1159-69.
 118. Gwee KA, Wee S, Wong ML, Png DJ. The prevalence, symptom characteristics, and impact of irritable bowel syndrome in an asian urban community. *Am.J.Gastroenterol.* 2004;99:924-31.
 119. Simren M, Abrahamsson H, Svedlund J, Björnsson ES. Quality of life in patients with irritable bowel syndrome seen in referral centers versus primary care: the impact of gender and predominant bowel pattern. *Scand.J.Gastroenterol.* 2001;36:545-52.
 120. Talley NJ, Boyce P, Jones M. Identification of distinct upper and lower gastrointestinal symptom groupings in an urban population. *Gut* 1998;42:690-5.
 121. Corney RH, Stanton R. Physical symptom severity, psychological and social dysfunction in a series of outpatients with irritable bowel syndrome. *J.Psychosom.Res.* 1990;34:483-91.
 122. Lee SY, Kim JH, Sung IK, Park HS, Jin CJ, Choe WH, Kwon SY, Lee CH, Choi KW. Irritable bowel syndrome is more common in women regardless of the menstrual phase: a Rome II-based survey. *J.Korean Med.Sci.* 2007;22:851-4.
 123. Lee OY, Mayer EA, Schmulson M, Chang L, Naliboff B. Gender-related differences in IBS symptoms. *Am.J.Gastroenterol.* 2001;96:2184-93.
 124. Talley NJ, Phillips SF, Melton LJ, Mulvihill C, Wiltgen C, Zinsmeister AR. Diagnostic value of the Manning criteria in irritable bowel syndrome. *Gut* 1990;31:77-81.
 125. Smith RC, Greenbaum DS, Vancouver JB, Henry RC, Reinhart MA, Greenbaum RB, Dean HA, Mayle JE. Gender differences in Manning criteria in the irritable bowel syndrome. *Gastroenterology* 1991;100:591-5.
 126. Bennett G, Talley NJ. Irritable bowel syndrome in the elderly. *Best Pract.Res.Clin.Gastroenterol.* 2002;16:63-76.
 127. Talley NJ, Zinsmeister AR, Melton LJ,3rd. Irritable bowel syndrome in a community: symptom subgroups, risk factors, and health care utilization. *Am.J.Epidemiol.* 1995;142:76-83.
 128. Minocha A, Johnson WD, Abell TL, Wigington WC. Prevalence, sociodemography, and quality of life of older versus younger patients with irritable bowel syndrome: a population-based study. *Dig.Dis.Sci.* 2006;51:446-53.
 129. Howell S, Talley NJ, Quine S, Poulton R. The irritable bowel syndrome has origins in the childhood socioeconomic environment. *Am.J.Gastroenterol.* 2004;99:1572-8.

130. Talley NJ, Howell S, Poulton R. The irritable bowel syndrome and psychiatric disorders in the community: is there a link? *Am.J.Gastroenterol.* 2001;96:1072-9.
131. Bommelaer G, Dorval E, Denis P, Czernichow P, Frexinos J, Pelc A, Slama A, El Hasnaoui A. Prevalence of irritable bowel syndrome in the French population according to the Rome I criteria. *Gastroenterol.Clin.Biol.* 2002;26:1118-23.
132. Dapoigny M, Bellanger J, Bonaz B, Bruley des Varannes S, Bueno L, Coffin B, Ducrotte P, Flourie B, Lemann M, Lepicard A, Reigenau O. Irritable bowel syndrome in France: a common, debilitating and costly disorder. *Eur.J.Gastroenterol.Hepatol.* 2004;16:995-1001.
133. Wilson S, Roberts L, Roalfe A, Bridge P, Singh S. Prevalence of irritable bowel syndrome: a community survey. *Br.J.Gen.Pract.* 2004;54:495-502.
134. Whitehead WE, Palsson O, Jones KR. Systematic review of the comorbidity of irritable bowel syndrome with other disorders: what are the causes and implications? *Gastroenterology* 2002;122:1140-56.
135. Vandvik PO, Lydersen S, Farup PG. Prevalence, comorbidity and impact of irritable bowel syndrome in Norway. *Scand.J.Gastroenterol.* 2006;41:650-6.
136. Lepine JP, Gastpar M, Mendlewicz J, Tylee A. Depression in the community: the first pan-European study DEPRES (Depression Research in European Society). *Int.Clin.Psychopharmacol.* 1997;12:19-29.
137. Kessler RC, McGonagle KA, Zhao S, Nelson CB, Hughes M, Eshleman S, Wittchen HU, Kendler KS. Lifetime and 12-month prevalence of DSM-III-R psychiatric disorders in the United States. Results from the National Comorbidity Survey. *Arch.Gen.Psychiatry* 1994;51:8-19.
138. Lindeman S, Hämäläinen J, Isometsä E, Kaprio J, Poikolainen K, Heikkinen M, Aro H. The 12-month prevalence and risk factors for major depressive episode in Finland: representative sample of 5993 adults. *Acta Psychiatr.Scand.* 2000;102:178-84.
139. Sartorius N, Ustun TB, Lecrubier Y, Wittchen HU. Depression comorbid with anxiety: results from the WHO study on psychological disorders in primary health care. *Br.J.Psychiatry Suppl.* 1996;(30):38-43.
140. Tollefson GD, Tollefson SL, Pederson M, Luxenberg M, Dunsmore G. Comorbid Irritable Bowel Syndrome in Patients with Generalized Anxiety and Major Depression. *Ann.Clin.Psychiatry* 1991;3:215-22.
141. Masand PS, Kaplan DS, Gupta S, Bhandary AN, Nasra GS, Kline MD, Margo KL. Major depression and irritable bowel syndrome: is there a relationship? *J.Clin.Psychiatry* 1995;56:363-7.
142. Katon W, Von Korff M, Lin E, Lipscomb P, Russo J, Wagner E, Polk E. Distressed high utilizers of medical care. DSM-III-R diagnoses and treatment needs. *Gen.Hosp.Psychiatry* 1990;12:355-62.

143. Kupfer DJ, Frank E. Comorbidity in depression. *Acta Psychiatr.Scand.Suppl.* 2003;(418):57-60.
144. Walker EA, Katon WJ, Roy-Byrne PP, Jemelka RP, Russo J. Histories of sexual victimization in patients with irritable bowel syndrome or inflammatory bowel disease. *Am.J.Psychiatry* 1993;150:1502-6.
145. Irwin C, Falsetti SA, Lydiard RB, Ballenger JC, Brock CD, Brenner W. Comorbidity of posttraumatic stress disorder and irritable bowel syndrome. *J.Clin.Psychiatry* 1996;57:576-8.
146. Lydiard RB, Fossey MD, Marsh W, Ballenger JC. Prevalence of psychiatric disorders in patients with irritable bowel syndrome. *Psychosomatics* 1993;34:229-34.
147. Henningsen P, Zimmermann T, Sattel H. Medically unexplained physical symptoms, anxiety, and depression: a meta-analytic review. *Psychosom.Med.* 2003;65:528-33.
148. Payne S. Sex, gender, and irritable bowel syndrome: making the connections. *Gend.Med.* 2004;1:18-28.
149. Mussell M, Kroenke K, Spitzer RL, Williams JB, Herzog W, Lowe B. Gastrointestinal symptoms in primary care: prevalence and association with depression and anxiety. *J.Psychosom.Res.* 2008;64:605-12.
150. Lydiard RB, Falsetti SA. Experience with anxiety and depression treatment studies: implications for designing irritable bowel syndrome clinical trials. *Am.J.Med.* 1999;107:65S-73S.
151. Riedl A, Schmidtmann M, Stengel A, Goebel M, Wisser AS, Klapp BF, Monnikes H. Somatic comorbidities of irritable bowel syndrome: a systematic analysis. *J.Psychosom.Res.* 2008;64:573-82.
152. Stanghellini V, Tosetti C, Barbara G, De Giorgio R, Cogliandro L, Cogliandro R, Corinaldesi R. Dyspeptic symptoms and gastric emptying in the irritable bowel syndrome. *Am.J.Gastroenterol.* 2002;97:2738-43.
153. Talley NJ, Dennis EH, Schettler-Duncan VA, Lacy BE, Olden KW, Crowell MD. Overlapping upper and lower gastrointestinal symptoms in irritable bowel syndrome patients with constipation or diarrhea. *Am.J.Gastroenterol.* 2003;98:2454-9.
154. van Bommel MJ, Numans ME, de Wit NJ, Stalman WA. Consultations and referrals for dyspepsia in general practice--a one year database survey. *Postgrad.Med.J.* 2001;77:514-8.
155. Nastaskin I, Mehdikhani E, Conklin J, Park S, Pimentel M. Studying the overlap between IBS and GERD: a systematic review of the literature. *Dig.Dis.Sci.* 2006;51:2113-20.
156. Donnelly VS, O'Herlihy C, Campbell DM, O'Connell PR. Postpartum fecal incontinence is more common in women with irritable bowel syndrome. *Dis.Colon Rectum* 1998;41:586-9.
157. Lee KJ, Kim JH, Cho SW. Relationship of underlying abnormalities in rectal sensitivity and compliance to distension with symptoms in irritable bowel syndrome. *Digestion* 2006;73:133-41.

158. Levy RL, Von Korff M, Whitehead WE, Stang P, Saunders K, Jhingran P, Barghout V, Feld AD. Costs of care for irritable bowel syndrome patients in a health maintenance organization. *Am.J.Gastroenterol.* 2001;96:3122-9.
159. Wolfe F, Ross K, Anderson J, Russell IJ, Hebert L. The prevalence and characteristics of fibromyalgia in the general population. *Arthritis Rheum.* 1995;38:19-28.
160. Jason LA, Richman JA, Rademaker AW, Jordan KM, Plioplys AV, Taylor RR, McCready W, Huang CF, Plioplys S. A community-based study of chronic fatigue syndrome. *Arch.Intern.Med.* 1999;159:2129-37.
161. Kennedy TM, Jones RH, Hungin AP, O'flanagan H, Kelly P. Irritable bowel syndrome, gastro-oesophageal reflux, and bronchial hyper-responsiveness in the general population. *Gut* 1998;43:770-4.
162. Ilnyckyj A, Graff LA, Blanchard JF, Bernstein CN. Therapeutic value of a gastroenterology consultation in irritable bowel syndrome. *Aliment.Pharmacol.Ther.* 2003;17:871-80.
163. Brody H. Placebo response, sustained partnership, and emotional resilience in practice. *J.Am.Board Fam.Pract.* 1997;10:72-4.
164. Thompson WG. Placebos: a review of the placebo response. *Am.J.Gastroenterol.* 2000;95:1637-43.
165. Bijkerk CJ, de Wit NJ, Stalman WA, Knottnerus JA, Hoes AW, Muris JW. Irritable bowel syndrome in primary care: the patients' and doctors' views on symptoms, etiology and management. *Can.J.Gastroenterol.* 2003;17:363,8; quiz 405-6.
166. van der Horst HE, van Dulmen AM, Schellevis FG, van Eijk JT, Fennis JF, Bleijenberg G. Do patients with irritable bowel syndrome in primary care really differ from outpatients with irritable bowel syndrome? *Gut* 1997;41:669-74.
167. Thompson WG, Heaton KW, Smyth GT, Smyth C. Irritable bowel syndrome in general practice: prevalence, characteristics, and referral. *Gut* 2000;46:78-82.
168. Whitehead WE, Levy RL, Von Korff M, Feld AD, Palsson OS, Turner M, Drossman DA. The usual medical care for irritable bowel syndrome. *Aliment.Pharmacol.Ther.* 2004;20:1305-15.
169. Colwell LJ, Prather CM, Phillips SF, Zinsmeister AR. Effects of an irritable bowel syndrome educational class on health-promoting behaviors and symptoms. *Am.J.Gastroenterol.* 1998;93:901-5.
170. Monsbakken KW, Vandvik PO, Farup PG. Perceived food intolerance in subjects with irritable bowel syndrome-- etiology, prevalence and consequences. *Eur.J.Clin.Nutr.* 2006;60:667-72.
171. Simren M, Mansson A, Langkilde AM, Svedlund J, Abrahamsson H, Bengtsson U, Björnsson ES. Food-related gastrointestinal symptoms in the irritable bowel syndrome. *Digestion* 2001;63:108-15.

172. Friedman G. Diet and the irritable bowel syndrome. *Gastroenterol.Clin.North Am.* 1991;20:313-24.
173. Mertz HR. Irritable bowel syndrome. *N.Engl.J.Med.* 2003;349:2136-46.
174. Suarez FL, Savaiano DA, Levitt MD. A comparison of symptoms after the consumption of milk or lactose-hydrolyzed milk by people with self-reported severe lactose intolerance. *N.Engl.J.Med.* 1995;333:1-4.
175. Vesa TH, Seppo LM, Marteau PR, Sahi T, Korpela R. Role of irritable bowel syndrome in subjective lactose intolerance. *Am.J.Clin.Nutr.* 1998;67:710-5.
176. Tolliver BA, Jackson MS, Jackson KL, Barnett ED, Chastang JF, DiPalma JA. Does lactose maldigestion really play a role in the irritable bowel? *J.Clin.Gastroenterol.* 1996;23:15-7.
177. Parker TJ, Woolner JT, Prevost AT, Tuffnell Q, Shorthouse M, Hunter JO. Irritable bowel syndrome: is the search for lactose intolerance justified? *Eur.J.Gastroenterol.Hepatol.* 2001;13:219-25.
178. Skoog SM, Bharucha AE. Dietary fructose and gastrointestinal symptoms: a review. *Am.J.Gastroenterol.* 2004;99:2046-50.
179. Shepherd SJ, Parker FC, Muir JG, Gibson PR. Dietary triggers of abdominal symptoms in patients with irritable bowel syndrome: randomized placebo-controlled evidence. *Clin.Gastroenterol.Hepatol.* 2008;6:765-71.
180. Dainese R, Galliani EA, De Lazzari F, Di Leo V, Naccarato R. Discrepancies between reported food intolerance and sensitization test findings in irritable bowel syndrome patients. *Am.J.Gastroenterol.* 1999;94:1892-7.
181. Brandt LJ, Chey WD, Foxx-Orenstein AE, Schiller LR, Schoenfeld PS, Spiegel BM, Talley NJ, Quigley EM. American College of Gastroenterology Task Force on Irritable Bowel Syndrome. *Am.J.Gastroenterol.* 2009;104:S1-S35.
182. Prior A, Whorwell PJ. Double blind study of ispaghula in irritable bowel syndrome. *Gut* 1987;28:1510-3.
183. Cann PA, Read NW, Holdsworth CD. What is the benefit of coarse wheat bran in patients with irritable bowel syndrome? *Gut* 1984;25:168-73.
184. Francis CY, Whorwell PJ. Bran and irritable bowel syndrome: time for reappraisal. *Lancet* 1994;344:39-40.
185. Bijkerk CJ, Muris JW, Knottnerus JA, Hoes AW, de Wit NJ. Systematic review: the role of different types of fibre in the treatment of irritable bowel syndrome. *Aliment.Pharmacol.Ther.* 2004;19:245-51.
186. Ford AC, Talley NJ, Spiegel BM, Foxx-Orenstein AE, Schiller L, Quigley EM, Moayyedi P. Effect of fibre, antispasmodics, and peppermint oil in the treatment of irritable bowel syndrome: systematic review and meta-analysis. *BMJ* 2008;337:a2313.

187. Lucey MR, Clark ML, Lowndes J, Dawson AM. Is bran efficacious in irritable bowel syndrome? A double blind placebo controlled crossover study. *Gut* 1987;28:221-5.
188. Bijkerk CJ, de Wit NJ, Muris JW, Whorwell PJ, Knottnerus JA, Hoes AW. Soluble or insoluble fibre in irritable bowel syndrome in primary care? Randomised placebo controlled trial. *BMJ* 2009;339:b3154.
189. Saxelin M, Tynkkynen S, Mattila-Sandholm T, de Vos WM. Probiotic and other functional microbes: from markets to mechanisms. *Curr.Opin.Biotechnol.* 2005;16:204-11.
190. Camilleri M. Probiotics and irritable bowel syndrome: rationale, mechanisms, and efficacy. *J.Clin.Gastroenterol.* 2008;42 Suppl 3 Pt 1:S123-5.
191. Lembo A. Irritable bowel syndrome medications side effects survey. *J.Clin.Gastroenterol.* 2004;38:776-81.
192. Brandt LJ, Bjorkman D, Fennerty MB, Locke GR, Olden K, Peterson W, Quigley E, Schoenfeld P, Schuster M, Talley N. Systematic review on the management of irritable bowel syndrome in North America. *Am.J.Gastroenterol.* 2002;97:S7-26.
193. Cann PA, Read NW, Holdsworth CD, Barends D. Role of loperamide and placebo in management of irritable bowel syndrome (IBS). *Dig.Dis.Sci.* 1984;29:239-47.
194. Efskind PS, Bernklev T, Vatn MH. A double-blind placebo-controlled trial with loperamide in irritable bowel syndrome. *Scand.J.Gastroenterol.* 1996;31:463-8.
195. Hovdenak N. Loperamide treatment of the irritable bowel syndrome. *Scand.J.Gastroenterol.Suppl.* 1987;130:81-4.
196. Lavo B, Stenstam M, Nielsen AL. Loperamide in treatment of irritable bowel syndrome--a double-blind placebo controlled study. *Scand.J.Gastroenterol.Suppl.* 1987;130:77-80.
197. Attar A, Lemann M, Ferguson A, Halphen M, Boutron MC, Flourie B, Alix E, Salmeron M, Guillemot F, Chaussade S, Menard AM, Moreau J, Naudin G, Barthet M. Comparison of a low dose polyethylene glycol electrolyte solution with lactulose for treatment of chronic constipation. *Gut* 1999;44:226-30.
198. Khoshoo V, Armstead C, Landry L. Effect of a laxative with and without tegaserod in adolescents with constipation predominant irritable bowel syndrome. *Aliment.Pharmacol.Ther.* 2006;23:191-6.
199. Jailwala J, Imperiale TF, Kroenke K. Pharmacologic treatment of the irritable bowel syndrome: a systematic review of randomized, controlled trials. *Ann.Intern.Med.* 2000;133:136-47.
200. Tack J, Fried M, Houghton LA, Spicak J, Fisher G. Systematic review: the efficacy of treatments for irritable bowel syndrome--a European perspective. *Aliment.Pharmacol.Ther.* 2006;24:183-205.
201. Cappello G, Spezzaferro M, Grossi L, Manzoli L, Marzio L. Peppermint oil (Mintoil®) in the treatment of irritable bowel syndrome: A prospective double blind placebo-controlled

- randomized trial. *Dig.Liver Dis.* 2007;39:530-6.
202. Liu JH, Chen GH, Yeh HZ, Huang CK, Poon SK. Enteric-coated peppermint-oil capsules in the treatment of irritable bowel syndrome: a prospective, randomized trial. *J.Gastroenterol.* 1997;32:765-8.
 203. McQuay HJ, Tramer M, Nye BA, Carroll D, Wiffen PJ, Moore RA. A systematic review of antidepressants in neuropathic pain. *Pain* 1996;68:217-27.
 204. Saarto T, Wiffen PJ. Antidepressants for neuropathic pain. *Cochrane Database Syst.Rev.* 2007;(4):CD005454.
 205. Hislop IG. Psychological significance of the irritable colon syndrome. *Gut* 1971;12:452-7.
 206. Clouse RE, Lustman PJ. Use of psychopharmacological agents for functional gastrointestinal disorders. *Gut* 2005;54:1332-41.
 207. Ford AC, Talley NJ, Schoenfeld PS, Quigley EM, Moayyedi P. Efficacy of antidepressants and psychological therapies in irritable bowel syndrome: systematic review and meta-analysis. *Gut* 2009;58:367-78.
 208. Drossman DA, Toner BB, Whitehead WE, Diamant NE, Dalton CB, Duncan S, Emmott S, Proffitt V, Akman D, Frusciante K, Le T, Meyer K, Bradshaw B, Mikula K, Morris CB, Blackman CJ, Hu Y, Jia H, Li JZ, Koch GG, Bangdiwala SI. Cognitive-behavioral therapy versus education and desipramine versus placebo for moderate to severe functional bowel disorders. *Gastroenterology* 2003;125:19-31.
 209. Vahedi H, Merat S, Momtahan S, Kazzazi AS, Ghaffari N, Olfati G, Malekzadeh R. Clinical trial: the effect of amitriptyline in patients with diarrhoea-predominant irritable bowel syndrome. *Aliment.Pharmacol.Ther.* 2008;27:678-84.
 210. O'Malley PG, Jackson JL, Santoro J, Tomkins G, Balden E, Kroenke K. Antidepressant therapy for unexplained symptoms and symptom syndromes. *J.Fam.Pract.* 1999;48:980-90.
 211. Creed F, Fernandes L, Guthrie E, Palmer S, Ratcliffe J, Read N, Rigby C, Thompson D, Tomenson B. The cost-effectiveness of psychotherapy and paroxetine for severe irritable bowel syndrome. *Gastroenterology* 2003;124:303-17.
 212. Tabas G, Beaves M, Wang J, Friday P, Mardini H, Arnold G. Paroxetine to treat irritable bowel syndrome not responding to high-fiber diet: a double-blind, placebo-controlled trial. *Am.J.Gastroenterol.* 2004;99:914-20.
 213. Kuiken SD, Tytgat GN, Boeckxstaens GE. The selective serotonin reuptake inhibitor fluoxetine does not change rectal sensitivity and symptoms in patients with irritable bowel syndrome: a double blind, randomized, placebo-controlled study. *Clin.Gastroenterol.Hepatol.* 2003;1:219-28.
 214. Vahedi H, Merat S, Rashidioon A, Ghoddoosi A, Malekzadeh R. The effect of fluoxetine in patients with pain and constipation-predominant irritable bowel syndrome: a double-blind randomized-controlled study. *Aliment.Pharmacol.Ther.* 2005;22:381-5.

215. Payne A, Blanchard EB. A controlled comparison of cognitive therapy and self-help support groups in the treatment of irritable bowel syndrome. *J.Consult.Clin.Psychol.* 1995;63:779-86.
216. Kennedy TM, Chalder T, McCrone P, Darnley S, Knapp M, Jones RH, Wessely S. Cognitive behavioural therapy in addition to antispasmodic therapy for irritable bowel syndrome in primary care: randomised controlled trial. *Health Technol.Assess.* 2006;10:iii,iv, ix-x, 1-67.
217. Galovski TE, Blanchard EB. The treatment of irritable bowel syndrome with hypnotherapy. *Appl.Psychophysiol.Biofeedback* 1998;23:219-32.
218. Simren M, Ringström G, Björnsson ES, Abrahamsson H. Treatment with hypnotherapy reduces the sensory and motor component of the gastrocolonic response in irritable bowel syndrome. *Psychosom.Med.* 2004;66:233-8.
219. Keefer L, Blanchard EB. The effects of relaxation response meditation on the symptoms of irritable bowel syndrome: results of a controlled treatment study. *Behav.Res.Ther.* 2001;39:801-11.
220. Boyce PM, Talley NJ, Balaam B, Koloski NA, Truman G. A randomized controlled trial of cognitive behavior therapy, relaxation training, and routine clinical care for the irritable bowel syndrome. *Am.J.Gastroenterol.* 2003;98:2209-18.
221. van der Veek PP, van Rood YR, Masclee AA. Clinical trial: short- and long-term benefit of relaxation training for irritable bowel syndrome. *Aliment.Pharmacol.Ther.* 2007;26:943-52.
222. Shaw G, Srivastava ED, Sadlier M, Swann P, James JY, Rhodes J. Stress management for irritable bowel syndrome: a controlled trial. *Digestion* 1991;50:36-42.
223. Blanchard EB, Schwarz SP, Suls JM, Gerardi MA, Scharff L, Greene B, Taylor AE, Berreman C, Malamood HS. Two controlled evaluations of multicomponent psychological treatment of irritable bowel syndrome. *Behav.Res.Ther.* 1992;30:175-89.
224. Heitkemper MM, Jarrett ME, Levy RL, Cain KC, Burr RL, Feld A, Barney P, Weisman P. Self-management for women with irritable bowel syndrome. *Clin.Gastroenterol.Hepatol.* 2004;2:585-96.
225. Guthrie E, Creed F, Dawson D, Tomenson B. A controlled trial of psychological treatment for the irritable bowel syndrome. *Gastroenterology* 1991;100:450-7.
226. Drossman DA, Whitehead WE, Camilleri M. Irritable bowel syndrome: a technical review for practice guideline development. *Gastroenterology* 1997;112:2120-37.
227. Shaheen NJ, Hansen RA, Morgan DR, Gangarosa LM, Ringel Y, Thiny MT, Russo MW, Sandler RS. The burden of gastrointestinal and liver diseases, 2006. *Am.J.Gastroenterol.* 2006;101:2128-38.
228. Thompson WG, Heaton KW. Functional bowel disorders in apparently healthy people. *Gastroenterology* 1980;79:283-8.
229. Talley NJ, Boyce PM, Jones M. Predictors of health care seeking for irritable bowel syndrome: a population based study. *Gut* 1997;41:394-8.

230. Koloski NA, Talley NJ, Huskic SS, Boyce PM. Predictors of conventional and alternative health care seeking for irritable bowel syndrome and functional dyspepsia. *Aliment.Pharmacol.Ther.* 2003;17:841-51.
231. Williams RE, Black CL, Kim HY, Andrews EB, Mangel AW, Buda JJ, Cook SF. Determinants of healthcare-seeking behaviour among subjects with irritable bowel syndrome. *Aliment.Pharmacol.Ther.* 2006;23:1667-75.
232. Gaburri M, Bassotti G, Bacci G, Cinti A, Bosso R, Ceccarelli P, Paolocci N, Pelli MA, Morelli A. Functional gut disorders and health care seeking behavior in an Italian non-patient population. *Recenti Prog.Med.* 1989;80:241-4.
233. Shih YC, Barghout VE, Sandler RS, Jhingran P, Sasane M, Cook S, Gibbons DC, Halpern M. Resource utilization associated with irritable bowel syndrome in the United States 1987-1997. *Dig.Dis.Sci.* 2002;47:1705-15.
234. Drossman DA, McKee DC, Sandler RS, Mitchell CM, Cramer EM, Lowman BC, Burger AL. Psychosocial factors in the irritable bowel syndrome. A multivariate study of patients and nonpatients with irritable bowel syndrome. *Gastroenterology* 1988;95:701-8.
235. Koloski NA, Talley NJ, Boyce PM. Predictors of health care seeking for irritable bowel syndrome and nonulcer dyspepsia: a critical review of the literature on symptom and psychosocial factors. *Am.J.Gastroenterol.* 2001;96:1340-9.
236. Ringström G, Abrahamsson H, Strid H, Simren M. Why do subjects with irritable bowel syndrome seek health care for their symptoms? *Scand.J.Gastroenterol.* 2007;42:1194-203.
237. Koloski NA, Talley NJ, Boyce PM. Epidemiology and health care seeking in the functional GI disorders: a population-based study. *Am.J.Gastroenterol.* 2002;97:2290-9.
238. Blewett A, Allison M, Calcraft B, Moore R, Jenkins P, Sullivan G. Psychiatric disorder and outcome in irritable bowel syndrome. *Psychosomatics* 1996;37:155-60.
239. Drossman DA, Li Z, Leserman J, Toomey TC, Hu YJ. Health status by gastrointestinal diagnosis and abuse history. *Gastroenterology* 1996;110:999-1007.
240. Prior A, Whorwell PJ. Gynaecological consultation in patients with the irritable bowel syndrome. *Gut* 1989;30:996-8.
241. Whitehead WE, Bosmajian LS. Behavioral medicine approaches to gastrointestinal disorders. *J.Consult.Clin.Psychol.* 1982;50:972-83.
242. Kettell J, Jones R, Lydeard S. Reasons for consultation in irritable bowel syndrome: symptoms and patient characteristics. *Br.J.Gen.Pract.* 1992;42:459-61.
243. Talley NJ, Gabriel SE, Harmsen WS, Zinsmeister AR, Evans RW. Medical costs in community subjects with irritable bowel syndrome. *Gastroenterology* 1995;109:1736-41.
244. Longstreth GF, Wilson A, Knight K, Wong J, Chiou CF, Barghout V, Frech F, Ofman JJ. Irritable bowel syndrome, health care use, and costs: a U.S. managed care perspective. *Am.J.Gastroenterol.* 2003;98:600-7.

245. Ricci J, Guerreiro T, Jones T, Schultes H, Volz A, Kessabi S. Irritable bowel syndrome: A comparison of the economic (direct and indirect) burden in four european countries. *Gut* 2002;51:A107-.
246. Le Pen C, Ruszniewski P, Gaudin AF, Amouretti M, Bommelaer G, Frexinos J, Poynard T, Maurel F, Priol G, Bertin C. The burden cost of French patients suffering from irritable bowel syndrome. *Scand.J.Gastroenterol.* 2004;39:336-43.
247. Badia X, Mearin F, Balboa A, Baro E, Caldwell E, Cucala M, Diaz-Rubio M, Fueyo A, Ponce J, Roset M, Talley NJ. Burden of illness in irritable bowel syndrome comparing Rome I and Rome II criteria. *Pharmacoeconomics* 2002;20:749-58.
248. Maxion-Bergemann S, Thielecke F, Abel F, Bergemann R. Costs of irritable bowel syndrome in the UK and US. *Pharmacoeconomics* 2006;24:21-37.
249. Hahn BA, Yan S, Strassels S. Impact of irritable bowel syndrome on quality of life and resource use in the United States and United Kingdom. *Digestion* 1999;60:77-81.
250. Sandler RS, Everhart JE, Donowitz M, Adams E, Cronin K, Goodman C, Gemmen E, Shah S, Avdic A, Rubin R. The burden of selected digestive diseases in the United States. *Gastroenterology* 2002;122:1500-11.
251. Leong SA, Barghout V, Birnbaum HG, Thibeault CE, Ben-Hamadi R, Frech F, Ofman JJ. The economic consequences of irritable bowel syndrome: a US employer perspective. *Arch.Intern.Med.* 2003;163:929-35.
252. Akehurst RL, Brazier JE, Mathers N, O'Keefe C, Kaltenthaler E, Morgan A, Platts M, Walters SJ. Health-related quality of life and cost impact of irritable bowel syndrome in a UK primary care setting. *Pharmacoeconomics* 2002;20:455-62.
253. Patel RP, Petitta A, Fogel R, Peterson E, Zarowitz BJ. The economic impact of irritable bowel syndrome in a managed care setting. *J.Clin.Gastroenterol.* 2002;35:14-20.
254. Inadomi JM, Fennerty MB, Bjorkman D. Systematic review: the economic impact of irritable bowel syndrome. *Aliment.Pharmacol.Ther.* 2003;18:671-82.
255. Lieberman DA, Holub J, Eisen G, Kraemer D, Morris CD. Utilization of colonoscopy in the United States: results from a national consortium. *Gastrointest.Endosc.* 2005;62:875-83.
256. Spiegel BM, Kanwal F, Naliboff B, Mayer E. The impact of somatization on the use of gastrointestinal health-care resources in patients with irritable bowel syndrome. *Am.J.Gastroenterol.* 2005;100:2262-73.
257. Yawn BP, Locke GR,3rd, Lydick E, Wollan PC, Bertram SL, Kurland MJ. Diagnosis and care of irritable bowel syndrome in a community-based population. *Am.J.Manag.Care* 2001;7:585-92.
258. Dean BB, Aguilar D, Barghout V, Kahler KH, Frech F, Groves D, Ofman JJ. Impairment in work productivity and health-related quality of life in patients with IBS. *Am.J.Manag.Care* 2005;11:S17-26.

259. Spiegel BM, Harris LA, Lucak SL, Mayer EA, Naliboff BD, Bolus RE, Esrailian E, Chey WD, Lembo A, Karsan HA, Tillisch K, Dulai GS, Strickland A, Chang L. Predictors of Work Productivity in Irritable Bowel Syndrome (IBS): Results from the Proof Cohort. *Gastroenterology* 2008;134:A-28.
260. Drossman D, Corazziari E, Delvaux M, Spiller RC, Talley NJ, Thompson WG, Whitehead WE, eds. *Rome III: The Functional Gastrointestinal Disorders*. Third edition. McLean, Virginia: Degnon Associates, 2006:837-43.
261. Beck AT, Beck RW. Screening depressed patients in family practice. A rapid technic. *Postgrad.Med.* 1972;52:81-5.
262. Beck AT, Rial WY, Rickels K. Short form of depression inventory: cross-validation. *Psychol.Rep.* 1974;34:1184-6.
263. Beck AT, Beamesderfer A. Assessment of depression: the depression inventory. *Mod.Probl.Pharmacopsychiatry* 1974;7:151-69.
264. Raitasalo R, Notkola V. Screening of mental health disorders among farmers and occupational health care. *J Soc Med* 1987;24:232-41.
265. Hänninen V, Aro H. Sex differences in coping and depression among young adults. *Soc.Sci.Med.* 1996;43:1453-60.
266. Kaltiala-Heino R, Rimpelä M, Rantanen P, Laippala P. Adolescent depression: the role of discontinuities in life course and social support. *J.Affect.Disord.* 2001;64:155-66.
267. Raitasalo R. Coping as the target of social policy. The Social Insurance Institution, Finland, *Studies in social security and health* 1. Helsinki 1995.
268. Aalto AM, Aro AR, Teperi J. RAND-36 terveyteen liittyvän elämänlaadun mittarina. 1st ed. Saarijärvi: STAKES Tutkimuksia 101; 1999.
269. Hujanen T. Terveystenhuollon yksikkökustannukset Suomessa vuonna 2001 (Unit costs in Finnish Health care in 2001). STAKES; 2003.
270. Drummond MF, Sculpher MJ, Torrance GW, O'Brien BJ, Stoddart GL editors. *Methods for the Economic Evaluation of Health Care Programmes*. Third ed. Toronto, Canada: Oxford University Press; 2005.
271. *Statistical Yearbook of Finland 2002*. Statistics Finland; 2003.
272. Fleiss JL editor. *Statistical Methods for Rates and Proportions*. 2nd ed. New York: John Wiley and Sons; 1981.
273. Barber JA, Thompson SG. Analysis of cost data in randomized trials: an application of the non-parametric bootstrap. *Stat.Med.* 2000;19:3219-36.
274. Furlanetto LM, Mendlowicz MV, Romildo Bueno J. The validity of the Beck Depression Inventory-Short Form as a screening and diagnostic instrument for moderate and severe depression in medical inpatients. *J.Affect.Disord.* 2005;86:87-91.

275. American College of Gastroenterology Task Force on Irritable Bowel Syndrome. An Evidence-Based Systematic Review on the Management of Irritable Bowel Syndrome. *Am.J.Gastroenterol.* 2009;104:S1-S35.
276. Drossman D, Corazziari E, Delvaux M, Spiller RC, Talley NJ, Thompson WG, Whitehead WE, eds. *Rome III: The Functional Gastrointestinal Disorders*. Third edition. McLean, Virginia: Degnon Associates, 2006:943.
277. Vandvik PO, Aabakken L, Farup PG. Diagnosing irritable bowel syndrome: poor agreement between general practitioners and the Rome II criteria. *Scand.J.Gastroenterol.* 2004;39:448-53.
278. Mearin F, Roset M, Badia X, Balboa A, Baro E, Ponce J, Diaz-Rubio M, Caldwell E, Cucala M, Fuego A, Talley NJ. Splitting irritable bowel syndrome: from original Rome to Rome II criteria. *Am.J.Gastroenterol.* 2004;99:122-30.
279. Pirkola SP, Isometsä E, Suvisaari J, Aro H, Joukamaa M, Poikolainen K, Koskinen S, Aromaa A, Lönnqvist JK. DSM-IV mood-, anxiety- and alcohol use disorders and their comorbidity in the Finnish general population--results from the Health 2000 Study. *Soc.Psychiatry Psychiatr.Epidemiol.* 2005;40:1-10.
280. Kaltiala-Heino R, Rimpelä M, Rantanen P, Laippala P. Finnish modification of the 13-item Beck Depression Inventory in screening an adolescent population for depressiveness and positive mood. *Nord.J.Psychiatry.* 1999;53:451-7.
281. Lydiard RB. Anxiety and the irritable bowel syndrome: psychiatric, medical, or both? *J.Clin.Psychiatry* 1997;58 Suppl 3:51,8; discussion 59-61.
282. Esler MD, Goulston KJ. Levels of anxiety in colonic disorders. *N.Engl.J.Med.* 1973;288:16-20.
283. Heefner JD, Wilder RM, Wilson ID. Irritable colon and depression. *Psychosomatics* 1978;19:540-7.
284. Mayer EA, Craske M, Naliboff BD. Depression, anxiety, and the gastrointestinal system. *J.Clin.Psychiatry* 2001;62 Suppl 8:28,36; discussion 37.
285. Nemeroff CB. The neurobiology of depression. *Sci.Am.* 1998;278:42-9.
286. Talley NJ, Dennis EH, Schettler-Duncan VA, Lacy BE, Olden KW, Crowell MD. Overlapping upper and lower gastrointestinal symptoms in irritable bowel syndrome patients with constipation or diarrhea. *Am.J.Gastroenterol.* 2003;98:2454-9.
287. Whorwell PJ, McCallum M, Creed FH, Roberts CT. Non-colonic features of irritable bowel syndrome. *Gut* 1986;27:37-40.
288. Maxton DG, Morris J, Whorwell PJ. More accurate diagnosis of irritable bowel syndrome by the use of 'non-colonic' symptomatology. *Gut* 1991;32:784-6.
289. Jones R, Latinovic R, Charlton J, Gulliford M. Physical and psychological co-morbidity in irritable bowel syndrome: a matched cohort study using the General Practice Research

Database. Aliment.Pharmacol.Ther. 2006;24:879-86.

290. White AM, Stevens WH, Upton AR, O'Byrne PM, Collins SM. Airway responsiveness to inhaled methacholine in patients with irritable bowel syndrome. *Gastroenterology* 1991;100:68-74.
291. Riccioni G, Della Vecchia R, Menna V, Staniscia T, Di Ilio C, Conti P, D'Orazio N. Irritable Bowel Syndrome and Bronchial Hyperresponsiveness: Is There a Link? *Digestion* 2004;69:185-9.
292. Jun DW, Lee OY, Yoon HJ, Lee HL, Yoon BC, Choi HS, Lee MH, Lee DH, Kee CS. Bronchial hyperresponsiveness in irritable bowel syndrome. *Dig.Dis.Sci.* 2005;50:1688-91.
293. Laatikainen T, Tapanainen H, Alfthan G, Salminen I, Sundvall J, Leiviskä J, Harald K, Jousilahti P, Salomaa P, Vartiainen E. Finnriski 2002. Publications of the National Public Health Institute. Helsinki 2003.
294. Hooper EW, Nycz GR, Regier DA, Goldberg ID, Jacobson A, Hankin J. Diagnosis of mental disorder in adults and increased use of health services in four outpatient settings. *Am.J.Psychiatry* 1980;137:207-10.
295. Lydeard S, Jones R. Factors affecting the decision to consult with dyspepsia: comparison of consulters and non-consulters. *J.R.Coll.Gen.Pract.* 1989;39:495-8.
296. Kwan AC, Hu WH, Chan YK, Yeung YW, Lai TS, Yuen H. Prevalence of irritable bowel syndrome in Hong Kong. *J.Gastroenterol.Hepatol.* 2002;17:1180-6.
297. El-Serag HB, Pilgrim P, Schoenfeld P. Systemic review: Natural history of irritable bowel syndrome. *Aliment.Pharmacol.Ther.* 2004;19:861-70.

1. Vastaaajan sukupuoli
☐ Nainen ☐ Mies
2. Vastaaajan ikä: vuotta**3. Siviilisääty**
☐ Naimaton
☐ Naimisissa
☐ Avoliitossa
☐ Leski
☐ Eronnut tai asumuserossa
4. Mikä on peruskoulutuksenne?
☐ Kansakoulu tai vähemmän
☐ Keskikoulu tai peruskoulu
☐ Osa lukiota
☐ Ylioppilas
5. Minkälaista työaikaa noudatatte?
☐ Säännöllinen päivätyö
☐ 2-vuorotyö
☐ 3-vuorotyö
☐ Yötyö
☐ Muu epäsäännöllinen työaika
☐ En ole töissä
6. Onko terveydentilanne yleisesti ottaen
☐ Erinomainen
☐ Varsin hyvä
☐ Hyvä
☐ Tyydyttävä
☐ Huono
7. Onko lääkäri todennut Teillä maitosokerin imeytymishäiriön eli laktoosi-intoleranssin?
☐ Ei ☐ Kyllä ☐ En osaa sanoa
8. Onko lääkäri todennut Teillä keliakia-sairauden?
☐ Ei ☐ Kyllä ☐ En osaa sanoa
9. Onko lääkäri todennut Teillä lapsuudessa (alle 15 -vuotiaana) seuraavia sairauksia:**a) Ruoka-aineallergia?**
☐ Ei ☐ Kyllä ☐ En osaa sanoa
b) Atooppinen ihottuma?
☐ Ei ☐ Kyllä ☐ En osaa sanoa
c) Astma?
☐ Ei ☐ Kyllä ☐ En osaa sanoa
d) Allerginen nuha?
☐ Ei ☐ Kyllä ☐ En osaa sanoa
10. Esiintyikö Teillä lapsuudessa (alle 15 -vuotiaana) muita enemmän vatsakipuja?
☐ Ei ☐ Kyllä ☐ En osaa sanoa
11. Onko umpilisäkkeenne leikattu?
☐ Ei ☐ Kyllä, milloin? Vuonna
12. Onko Teille tehty sappileikkaus?
☐ Ei ☐ Kyllä, milloin? Vuonna
13. Onko Teille tehty jokin muu suoliston tai vatsan alueen leikkaus?
☐ Ei ☐ Kyllä, milloin? Vuonna
Mikä leikkaus?

14. Onko Teille tehty seuraavia mahasuolikanavan tutkimuksia:**a) Mahalaukun tähystys?**
☐ Ei ☐ Kyllä, viimeksi vuonna
b) Paksusuolen tähystys?
☐ Ei ☐ Kyllä, viimeksi vuonna
15. Onko Teillä jokin pitkäaikaissairaus?
☐ Ei ☐ Kyllä, mikä?

16. Onko Teillä säännöllinen lääkitys?
☐ Ei ☐ Kyllä, mikä?

Listatkaa tähän kaikki säännöllisesti käyttämäenne lääkkeet, myös käsikauppalääkkeet:

17. Kuinka usein Teillä on ollut vatsakipua tai vatsavaivaa (vatsan seudun epämiellyttävää tunnetta) viimeksi kuluneen vuoden aikana?

- ☐ Harvoin tai ei ollenkaan
☐ Toisinaan (useammin kuin yhtenä päivänä kymmenestä)
☐ Usein (useammin kuin yhtenä päivänä neljästä)
☐ Hyvin usein (vähintään joka toinen päivä)
☐ Jatkuvasti (päivittäin)

Mikäli vastasitte "harvoin tai ei ollenkaan", voitte siirtyä suoraan kysymykseen 30, muussa tapauksessa vastatkaa vielä seuraaviin kysymyksiin:

18. Kuinka usein viimeksi kuluneen vuoden aikana vatsakipu tai vatsavaiva häiritsi päivittäisiä askareitanne?

- ☐ Harvoin tai ei ollenkaan
☐ Toisinaan (useammin kuin yhtenä päivänä kymmenestä)
☐ Usein (useammin kuin yhtenä päivänä neljästä)
☐ Hyvin usein (vähintään joka toinen päivä)
☐ Jatkuvasti (päivittäin)

19. Kuinka voimakasta edellä mainittu vatsaoire on yleensä ollut viimeisen vuoden aikana?

- ☐ Lievää (voi helposti jäädä huomaamatta, ellei sitä ajattele)
☐ Kohtalaista (oire on olemassa, mutta ei häiritse päivän toimintoja)
☐ Voimakasta (oire häiritsee päivän toimintoja)
☐ Erittäin voimakasta (oire häiritsee merkittävästi päivän toimintoja)

20. Helpottuiko edellä mainittu vatsakipu tai vatsavaiva ulostamisen jälkeen?

- ☐ Harvoin tai ei ollenkaan
☐ Toisinaan (useammin kuin yhtenä kertana kymmenestä)
☐ Usein (useammin kuin yhtenä kertana neljästä)
☐ Hyvin usein (vähintään joka toinen kerta)
☐ Lähes aina

21. Muuttuiko suolentoimintanne (ulostamistiheys) tavallista tiheämmäksi edellä mainitun vatsakivun tai -vaivan alkaessa?

- ☐ Harvoin tai ei ollenkaan
☐ Toisinaan (useammin kuin yhtenä kertana kymmenestä)
☐ Usein (useammin kuin yhtenä kertana neljästä)
☐ Hyvin usein (vähintään joka toinen kerta)
☐ Lähes aina

22. Muuttuiko suolentoimintanne tavallista harvemmaksi edellä mainitun vatsakivun tai -vaivan alkaessa?

- ☐ Harvoin tai ei ollenkaan
☐ Toisinaan (useammin kuin yhtenä kertana kymmenestä)
☐ Usein (useammin kuin yhtenä kertana neljästä)
☐ Hyvin usein (vähintään joka toinen kerta)
☐ Lähes aina

23. Muuttuiko ulosteenne koostumus tavallista löysemmäksi edellä mainitun vatsakivun tai -vaivan alkaessa?

- ☐ Harvoin tai ei ollenkaan
☐ Toisinaan (useammin kuin yhtenä kertana kymmenestä)
☐ Usein (useammin kuin yhtenä kertana neljästä)
☐ Hyvin usein (vähintään joka toinen kerta)
☐ Lähes aina

24. Muuttuiko ulosteenne koostumus tavallista kovemaksi edellä mainitun vatsakivun tai -vaivan alkaessa?

- ☐ Harvoin tai ei ollenkaan
☐ Toisinaan (useammin kuin yhtenä kertana kymmenestä)
☐ Usein (useammin kuin yhtenä kertana neljästä)
☐ Hyvin usein (vähintään joka toinen kerta)
☐ Lähes aina

25. Mitkä tekijät pahentavat vatsaoireitanne?**a) Ruokailu?**

- ☐ Ei pahenna ☐ Pahentaa joskus oireita ☐ Pahentaa usein oireita
☐ Pahentaa aina oireita ☐ En osaa sanoa

b) Maito tai maitotuotteet?

- ☐ Eivät pahenna ☐ Pahentavat joskus oireita ☐ Pahentavat usein oireita
☐ Pahentavat aina oireita ☐ En osaa sanoa

Mitä oireita saatte maitotuotteista? _____

Mitkä maitotuotteet aiheuttavat oireita? _____

c) Viljatuotteet?

- ☐ Eivät pahenna ☐ Pahentavat joskus oireita ☐ Pahentavat usein oireita
☐ Pahentavat aina oireita ☐ En osaa sanoa

Mitä oireita saatte viljatuotteista? _____

Mitkä viljatuotteet aiheuttavat oireita? _____

d) Jotkin muut tekijät tai ruoka-aineet? Mitkä? _____

Minkälaisia oireita ne aiheuttavat? _____

26. Onko Teillä ollut viimeksi kuluneen vuoden aikana usein vatsakipua tai vatsavaivaa (vatsan seudun epämiellyttävää tunnetta)? Usein tarkoittaa tässä yhteydessä oireen esiintymistä vähintään kerran viikossa ainakin 12 eri viikkona viimeisen vuoden aikana.

- ☐ Ei ☐ Kyllä

27. Kuinka usein viimeksi kuluneen vuoden aikana Teillä on ollut ylävatsalla (navan yläpuolella) tuntuvaa kipua? Närästystä eli rintalastan takana tuntuvaa polttelua ei lasketa mukaan.

- ☐ Harvoin tai ei ollenkaan
☐ Toisinaan (useammin kuin yhtenä päivänä kymmenestä)
☐ Usein (useammin kuin yhtenä päivänä neljästä)
☐ Hyvin usein (vähintään joka toinen päivä)
☐ Jatkuvasti tai lähes jatkuvasti

28. Kuinka usein viimeisen vuoden aikana Teillä on ollut ylävatsalla tuntuvaa muuta epämukavuuden tunnetta, esim. pahoinvointia, ylävatsan turvotusta tai ruokailun yhteydessä varhaista vatsan täyttymisen tunnetta?

- ☐ Harvoin tai ei ollenkaan
☐ Toisinaan (useammin kuin yhtenä päivänä kymmenestä)
☐ Usein (useammin kuin yhtenä päivänä neljästä)
☐ Hyvin usein (vähintään joka toinen päivä)
☐ Jatkuvasti tai lähes jatkuvasti

Jos vastasitte "harvoin tai ei ollenkaan", voitte siirtyä kysymykseen 30, muussa tapauksessa vastatkaa vielä seuraavaan:

29. Kuinka voimakasta edellä mainittu ylävatsaoire on yleensä ollut viimeisen vuoden aikana?

- ☐ Lievää (voi helposti jäädä huomaamatta, ellei sitä ajattele)
☐ Kohtalaista (oire on olemassa, mutta ei häiritse päivän toimintoja)
☐ Voimakasta (oire häiritsee päivän toimintoja)
☐ Erittäin voimakasta (oire häiritsee merkittävästi päivän toimintoja)

30. Kuinka usein viimeksi kuluneen vuoden aikana Teillä oli enemmän kuin kolme ulostuskertaa päivässä?

- ☐ Harvoin tai ei ollenkaan
- ☐ Toisinaan (useammin kuin yhtenä päivänä kymmenestä)
- ☐ Usein (useammin kuin yhtenä päivänä neljästä)
- ☐ Hyvin usein (vähintään joka toinen päivä)
- ☐ Jatkuvasti tai lähes jatkuvasti

31. Kuinka usein viimeksi kuluneen vuoden aikana Teillä oli vähemmän kuin kolme ulostuskertaa viikossa?

- ☐ Harvoin tai ei ollenkaan
- ☐ Toisinaan (useammin kuin yhtenä viikkona kymmenestä)
- ☐ Usein (useammin kuin yhtenä viikkona neljästä)
- ☐ Hyvin usein (vähintään joka toinen viikko)
- ☐ Jatkuvasti tai lähes jatkuvasti

32. Kuinka usein viimeksi kuluneen vuoden aikana ulosteenne oli kovaa, kokkaremaista tai papanamaista?

- ☐ Harvoin tai ei ollenkaan
- ☐ Toisinaan (useammin kuin yhtenä kertana kymmenestä)
- ☐ Usein (useammin kuin yhtenä kertana neljästä)
- ☐ Hyvin usein (vähintään joka toinen kerta)
- ☐ Jatkuvasti tai lähes jatkuvasti

33. Kuinka usein viimeksi kuluneen vuoden aikana ulosteenne oli löysää tai vetistä?

- ☐ Harvoin tai ei ollenkaan
- ☐ Toisinaan (useammin kuin yhtenä kertana kymmenestä)
- ☐ Usein (useammin kuin yhtenä kertana neljästä)
- ☐ Hyvin usein (vähintään joka toinen kerta)
- ☐ Jatkuvasti tai lähes jatkuvasti

34. Kuinka usein viimeksi kuluneen vuoden aikana Teille jäi ulostamistapahtuman jälkeen tunne, että peräsuolenne ei tyhjentynyt kunnolla?

- ☐ Harvoin tai ei ollenkaan
- ☐ Toisinaan (useammin kuin yhtenä kertana kymmenestä)
- ☐ Usein (useammin kuin yhtenä kertana neljästä)
- ☐ Hyvin usein (vähintään joka toinen kerta)
- ☐ Jatkuvasti tai lähes jatkuvasti

35. Kuinka usein viimeksi kuluneen vuoden aikana Teidän oli ponnisteltava runsaasti saadaksenne peräsuolenne tyhjentymään?

- ☐ Harvoin tai ei ollenkaan
- ☐ Toisinaan (useammin kuin yhtenä kertana kymmenestä)
- ☐ Usein (useammin kuin yhtenä kertana neljästä)
- ☐ Hyvin usein (vähintään joka toinen kerta)
- ☐ Aina tai lähes aina

36. Kuinka usein viimeksi kuluneen vuoden aikana Teillä on ollut pakottava ulostamisen tarve, jonka vuoksi Teidän on täytynyt kiirehtiä wc:hen?

- ☐ Harvoin tai ei ollenkaan
- ☐ Toisinaan (useammin kuin yhtenä kertana kymmenestä)
- ☐ Usein (useammin kuin yhtenä kertana neljästä)
- ☐ Hyvin usein (vähintään joka toinen kerta)
- ☐ Aina tai lähes aina

37. Kuinka usein viimeksi kuluneen vuoden aikana olette huomannut limaa ulosteissanne?

- ☐ Harvoin tai ei ollenkaan
- ☐ Toisinaan (useammin kuin yhtenä kertana kymmenestä)
- ☐ Usein (useammin kuin yhtenä kertana neljästä)
- ☐ Hyvin usein (vähintään joka toinen kerta)
- ☐ Aina tai lähes aina

38. Kuinka usein viimeksi kuluneen vuoden aikana Teillä on ollut vatsan turvotusta?

- ☐ Harvoin tai ei ollenkaan
- ☐ Toisinaan (useammin kuin yhtenä päivänä kymmenestä)
- ☐ Usein (useammin kuin yhtenä päivänä neljästä)
- ☐ Hyvin usein (vähintään joka toinen päivä)
- ☐ Jatkuvasti tai lähes jatkuvasti

39. Kuinka monta kertaa viimeisen vuoden aikana olette käynyt lääkärissä vatsavaivojen vuoksi?

- ☐ En kertaakaan
- ☐ 1-2 kertaa
- ☐ 3-5 kertaa
- ☐ 6-10 kertaa
- ☐ Yli 10 kertaa

40. Kuinka monta kertaa viimeisen vuoden aikana olette käynyt lääkärin vastaanotolla muiden syiden kuin vatsavaivojen vuoksi?

- ☐ En kertaakaan
- ☐ 1-2 kertaa
- ☐ 3-5 kertaa
- ☐ 6-10 kertaa
- ☐ Yli 10 kertaa

41. Kuinka monta päivää olette ollut vatsavaivojen takia poissa töistä viimeisen vuoden aikana?

- ☐ 0 päivää
- ☐ 1-3 päivää
- ☐ 4-6 päivää
- ☐ 7-20 päivää
- ☐ Yli 20 päivää
- ☐ En ole ollut työssä viimeisen vuoden aikana

42. Kuinka usein olette viimeisen vuoden aikana käyttänyt vatsaoireiden vuoksi lääkärin määräämiä lääkkeitä?

- ☐ En lainkaan
- ☐ Harvemmin kuin kerran kuukaudessa
- ☐ Keskimäärin 1-3 kertaa kuukaudessa
- ☐ Keskimäärin 1-3 kertaa viikossa
- ☐ Päivittäin tai lähes päivittäin

43. Kuinka usein olette viimeisen vuoden aikana käyttänyt vatsaoireiden vuoksi reseptivapaita käsikauppalääkkeitä?

- ☐ En lainkaan
- ☐ Harvemmin kuin kerran kuukaudessa
- ☐ Keskimäärin 1-3 kertaa kuukaudessa
- ☐ Keskimäärin 1-3 kertaa viikossa
- ☐ Päivittäin tai lähes päivittäin

44. Kuinka usein Teillä esiintyy seuraavia oireita?**a) Päänsärkyä?**

- ☐ Harvemmin kuin kerran kuukaudessa
- ☐ 1-3 kertaa kuukaudessa
- ☐ 1-3 kertaa viikossa
- ☐ Useammin kuin 3:sti viikossa

b) Selkäkipua?

- ☐ Harvemmin kuin kerran kuukaudessa
- ☐ 1-3 kertaa kuukaudessa
- ☐ 1-3 kertaa viikossa
- ☐ Useammin kuin 3:sti viikossa

Lopuksi esitämme eräitä kysymyksiä, jotka käsittelevät mielialan erilaisia piirteitä. Vastatkaa kuhunkin kysymykseen merkitsemällä rasti siihen kysymyksen jäljessä olevaan ruutuun, joka parhaiten kuvaa Teidän tilannettanne tällä hetkellä. Valitkaa kustakin kysymyksestä vain yksi vaihtoehto älkääkä jättäkö yhtään kysymystä väliin.

45. Minkälainen Teidän mielialanne on?

- ☐ Mielialani on melko valoisa ja hyvä
- ☐ En ole alakuloinen tai surullinen
- ☐ Tunnen itseni alakuloiseksi tai surulliseksi
- ☐ Olen alakuloinen jatkuvasti enkä pääse siitä
- ☐ Olen niin masentunut ja alavireinen, etten kestä enää

46. Miten suhtaudutte tulevaisuuteen?

- ☐ Suhtaudun tulevaisuuteeni toiveikkaasti
- ☐ En suhtaudu tulevaisuuteeni toivottomasti
- ☐ Tulevaisuus tuntuu minusta melko masentavalta
- ☐ Minusta tuntuu, ettei minulla ole tulevaisuudelta mitään odotettavaa
- ☐ Tulevaisuus tuntuu minusta toivottomalta, enkä jaksa uskoa, että asiat muuttuisivat parempaan päin

47. Miten katsotte elämänne sujuneen?

- ☐ Olen elämässäni onnistunut huomattavan usein
- ☐ En tunne epäonnistuneeni elämässäni
- ☐ Minusta tuntuu, että olen epäonnistunut pyrkimyksissäni tavallista useammin
- ☐ Elämäni on tähän saakka ollut vain sarja epäonnistumisia
- ☐ Tunnen epäonnistuneeni täydellisesti ihmisenä

48. Miten tyytyväiseksi tai tyytymättömäksi tunnette itsenne?

- ☐ Olen varsin tyytyväinen elämäni
- ☐ En ole erityisen tyytymätön
- ☐ En nauti asioista samalla tavalla kuin ennen
- ☐ Minusta tuntuu, etten saa tyydytystä juuri mistään
- ☐ Olen täysin tyytymätön kaikkeen

49. Minkälaisena pidätte itseänne?

- ☐ Tunnen itseni melko hyväksi
- ☐ En tunne itseäni huonoksi ja arvottomaksi
- ☐ Tunnen itseni huonoksi ja arvottomaksi melko usein
- ☐ Nykyään tunnen itseni arvottomaksi melkein aina
- ☐ Olen kerta kaikkiaan arvoton ja huono

50. Onko Teillä pettymyksen tunteita?

- ☐ Olen tyytyväinen itseeni ja suorituksiini
- ☐ En ole pettynyt itсени suhteen
- ☐ Olen pettynyt itсени suhteen
- ☐ Minua inhottaa oma itсени
- ☐ Vihaan itseäni

51. Onko Teillä itsenne vahingoittamiseen liittyviä ajatuksia?

- ☐ Minulla ei ole koskaan ollut itsemurha-ajatuksia
- ☐ En ajattele enkä halua vahingoittaa itseäni
- ☐ Minusta tuntuu, että olisi parempi, jos olisin kuollut
- ☐ Minulla on tarkat suunnitelmat itsemurhasta
- ☐ Tekisin itsemurhan, jos siihen olisi mahdollisuus

52. Miten suhtaudutte vieraitten ihmisten tapaamiseen?

- ☐ Pidän ihmisten tapaamisesta ja juttelemisesta
- ☐ En ole menettänyt kiinnostustani muihin ihmisiin
- ☐ Toiset ihmiset eivät enää kiinnosta minua niin paljon kuin ennen
- ☐ Olen melkein kokonaan menettänyt mielenkiintoni ja tunteeni toisia ihmisiä kohtaan
- ☐ Olen menettänyt mielenkiintoni muihin ihmisiin, enkä välitä heistä lainkaan

53. Miten koette päätösten tekemisen?

- ☐ Erilaisten päätösten tekeminen on minulle helppoa
- ☐ Pystyn tekemään päätöksiä samoin kuin ennenkin
- ☐ Varmuuteni on vähentynyt ja yritän lykätä päätösten tekoa
- ☐ Minulla on suuria vaikeuksia päätösten teossa
- ☐ En pysty enää lainkaan tekemään ratkaisuja ja päätöksiä

54. Minkälaisena pidätte olemustanne ja ulkonäköänne?

- ☐ Olen melko tyytyväinen ulkonäkööni ja olemukseeni
- ☐ Ulkonäössäni ei ole minua haittaavia piirteitä
- ☐ Olen huolissani siitä, että näytän epämiellyttävältä
- ☐ Minusta tuntuu, että näytän rumalta
- ☐ Olen varma, että näytän rumalta ja vastenmieliseltä

55. Minkälaista Teidän nukkumisenne on?

- ☐ Minulla ei ole nukkumisessa minkäänlaisia vaikeuksia
- ☐ Nukun yhtä hyvin kuin ennenkin
- ☐ Herätessäni aamuisin olen paljon väsyneempi kuin ennen
- ☐ Minua häiritsee unettomuus
- ☐ Kärsin unettomuudesta, nukahtamisvaikeuksista tai liian aikaisin kesken unien heräämisistä

56. Tunnetteko väsymystä tai uupumusta?

- ☐ Väsyminen on minulle lähes täysin vierasta
- ☐ En väsy helpommin kuin tavallisestikaan
- ☐ Väsyn nopeammin kuin ennen
- ☐ Vähäinenkin työ väsyttää ja uuvuttaa minua
- ☐ Olen liian väsynyt tehdäkseeni mitään

57. Minkälainen ruokahalunne on?

- ☐ Ruokahalussani ei ole mitään hankaluuksia
- ☐ Ruokahaluni on ennallaan
- ☐ Ruokahaluni on huonompi kuin ennen
- ☐ Ruokahaluni on nyt paljon huonompi kuin ennen
- ☐ Minulla ei ole enää lainkaan ruokahalua

58. Oletteko ahdistunut tai jännittynyt?

- ☐ Pidän itseäni melko hyvähermoisena enkä ahdistu kovinkaan helposti
- ☐ En tunne itseäni ahdistuneeksi tai "huonohermoiseksi"
- ☐ Ahdistun ja jännityn melko helposti
- ☐ Tulen erityisen helposti tuskaiseksi, ahdistuneeksi tai jännittyneeksi
- ☐ Tunnen itseni jatkuvasti ahdistuneeksi ja tuskaiseksi kuin hermoni olisivat "loppuun kuluneet"

Tarkistakaa vielä, että olette vastannut kaikkiin kysymyksiin ja palauttakaa kaavake oheisessa palautuskuoressa. KIITOS VAIVANNÄÖSTÄNNE!

1. Kuinka kauan Teillä on ollut vatsavaivoja?

- ☐ Alle vuoden ajan ☐ 1-3 vuotta ☐ 3-10 vuotta ☐ Yli 10 vuotta

2. Ovatko vatsaoireenne viimeisen 3 kuukauden aikana pysyneet ennallaan, lisääntyneet vai vähentyneet?

- ☐ Pysyneet ennallaan ☐ Lisääntyneet ☐ Vähentyneet

3. Luettelemme seuraavaksi 5 erilaista vatsaoiretta. Nimetkää näistä pahin oireenne laittamalla numero 1 oireen vieressä olevaan ruutuun, toiseksi pahin numerolla 2 jne. Mikäli kärsitte jostakin muusta vatsaoireesta, kirjoittakaa se vapaaseen tilaan ja merkitkää sen viereen sopivaksi katsomanne numero oireen hankaluuden mukaan.

Vatsakipu	<input type="text"/>
Vatsan turvotus.....	<input type="text"/>
Ilmavaivat.....	<input type="text"/>
Ripuli.....	<input type="text"/>
Ummetus.....	<input type="text"/>
Jokin muu oire, mikä?	<input type="text"/>

4. Kuinka usein Teillä on ollut vatsakipua tai vatsavaivaa (vatsan seudun epämiellyttävää tunnetta) viimeksi kuluneen 3 kuukauden aikana?

- ☐ Harvoin tai ei ollenkaan
☐ Toisinaan (useammin kuin yhtenä päivänä kymmenestä)
☐ Usein (useammin kuin yhtenä päivänä neljästä)
☐ Hyvin usein (vähintään joka toinen päivä)
☐ Jatkuvasti (päivittäin)


Mikäli vastasitte "harvoin tai ei ollenkaan", voitte siirtyä seuraavalle sivulle kysymykseen 7, muussa tapauksessa vastatkaa vielä seuraaviin kysymyksiin:

5. Millä alueella vatsakipunne tai -vaivanne tuntuu? Valitkaa oheisen kaavakuvan mukaisesti tilannettanne parhaiten kuvaava alue A,B,C tai D. Voitte valita myös useamman kuin yhden kirjaimen, mikäli oireenne tuntuu laajemmalla alueella. Merkitkää vastauksenne rastilla (X) asianomaisten kirjainten kohdalla oleviin ruutuihin.

- ☐ A
☐ B
☐ C
☐ D

6. Mikäli valitsitte edelliseen kysymykseen useamman kuin yhden alueen, pyydämme Teitä vielä arvioimaan, millä näistä alueista oleva kipu tai vaiva häiritsee Teitä eniten? Valitkaa nyt vain yksi alue yllä olevasta kaavakuvasta.

- ☐ A
☐ B
☐ C
☐ D
☐ En osaa sanoa



7. Kuinka monta kertaa Teille on tehty vatsavaivojenne vuoksi seuraavia tutkimuksia?

Paksusuolen tähystys?

☐ Ei kertaakaan ☐ Kerran ☐ 2 kertaa ☐ 3 kertaa ☐ Yli 3 kertaa

Paksusuolen röntgenkuvaus?

☐ Ei kertaakaan ☐ Kerran ☐ 2 kertaa ☐ 3 kertaa ☐ Yli 3 kertaa

Mahalaukun tähystys?

☐ Ei kertaakaan ☐ Kerran ☐ 2 kertaa ☐ 3 kertaa ☐ Yli 3 kertaa

Vatsan alueen ultraäänitutkimus?

☐ Ei kertaakaan ☐ Kerran ☐ 2 kertaa ☐ 3 kertaa ☐ Yli 3 kertaa

8. Oletteko käynyt lääkärissä vatsavaivojenne takia viimeisen vuoden aikana?

☐ Kyllä ☐ En

Mikäli vastasitte "En", voitte siirtyä seuraavalle sivulle kysymykseen 14, mikäli vastasitte "Kyllä", jatkakaa seuraavasta kysymyksestä:

9. Kuinka monta kertaa viimeisen vuoden aikana olette käynyt vatsavaivojenne takia terveyskeskuslääkärin vastaanotolla?

☐ En kertaakaan ☐ Kerran ☐ 2 kertaa ☐ 3 kertaa ☐ 4-6 kertaa ☐ yli 6 kertaa

10. Kuinka monta kertaa viimeisen vuoden aikana olette käynyt vatsavaivojenne takia työterveyslääkärin vastaanotolla?

☐ En kertaakaan ☐ Kerran ☐ 2 kertaa ☐ 3 kertaa ☐ 4-6 kertaa ☐ yli 6 kertaa

11. Kuinka monta kertaa viimeisen vuoden aikana olette käynyt vatsavaivojenne takia sairaalan ajanvarauspoliklinikalla (sisätautien tai kirurgian poliklinikalla)?

☐ En kertaakaan ☐ Kerran ☐ 2 kertaa ☐ 3 kertaa ☐ 4-6 kertaa ☐ yli 6 kertaa

12. Kuinka monta kertaa viimeisen vuoden aikana olette käynyt vatsavaivojenne takia sairaalan päivystyspoliklinikalla (ensiapupoliklinikalla)?

☐ En kertaakaan ☐ Kerran ☐ 2 kertaa ☐ 3 kertaa ☐ 4-6 kertaa ☐ yli 6 kertaa

13. Kuinka monta kertaa viimeisen vuoden aikana olette käynyt vatsavaivojenne takia yksityislääkärin vastaanotolla? (Tässä ei tarkoiteta käyntejä yksityisellä lääkäriasemalla työterveyslääkärin luona)

☐ En kertaakaan ☐ Kerran ☐ 2 kertaa ☐ 3 kertaa ☐ 4-6 kertaa ☐ yli 6 kertaa

14. Kuinka usein viimeksi kuluneen 3 kuukauden aikana vatsaoireenne on haitannut Teidän työntekoanne töissä ollessanne?

- ☐ Ei ollenkaan
- ☐ Harvemmin kuin 2 päivänä kuukaudessa
- ☐ 2-3 päivänä kuukaudessa
- ☐ 1-2 päivänä viikossa
- ☐ 3 päivänä viikossa tai useammin
- ☐ En ole ollut töissä viimeisten 3 kuukauden aikana

15. Kuinka usein viimeksi kuluneen 3 kuukauden aikana olette käyttänyt vatsavaivojenne takia jotain seuraavista reseptivapaista "näristyslääkkeistä": Balancid, Gaviscon, Link, Magnesiamaito, Novaluzid, PeeHoo, PH maxi, Rennie, Alsucral, Antepsin?

- ☐ En kertaakaan
- ☐ Harvemmin kuin 2 kertaa kuukaudessa
- ☐ Keskimäärin 2-3 kertaa kuukaudessa
- ☐ Keskimäärin 1-2 kertaa viikossa
- ☐ Keskimäärin 3-5 kertaa viikossa
- ☐ Päivittäin tai lähes päivittäin

16. Kuinka usein viimeksi kuluneen 3 kuukauden aikana olette käyttänyt vatsavaivojenne takia jotain seuraavista mahan suolahapon erityistä vähentävistä lääkkeistä: Esofex, Ranicur, Ranil, Ranimex, Ranitidine, Ranitidin, Ranixal, Zantac, Pepcid, Pepcidin, Nizax?

- ☐ En kertaakaan
- ☐ Harvemmin kuin 2 kertaa kuukaudessa
- ☐ Keskimäärin 2-3 kertaa kuukaudessa
- ☐ Keskimäärin 1-2 kertaa viikossa
- ☐ Keskimäärin 3-5 kertaa viikossa
- ☐ Päivittäin tai lähes päivittäin

17. Kuinka usein viimeksi kuluneen 3 kuukauden aikana olette käyttänyt vatsavaivojenne takia jotain seuraavista mahan suolahapon erityistä vähentävistä lääkkeistä: Lanzo, Losec, Nexium, Pariet, Somac?

- ☐ En kertaakaan
- ☐ Harvemmin kuin 2 kertaa kuukaudessa
- ☐ Keskimäärin 2-3 kertaa kuukaudessa
- ☐ Keskimäärin 1-2 kertaa viikossa
- ☐ Keskimäärin 3-5 kertaa viikossa
- ☐ Päivittäin tai lähes päivittäin

18. Kuinka usein viimeksi kuluneen 3 kuukauden aikana olette käyttänyt vatsavaivojenne takia reseptivapaata kipulääkettä, kuten Aspirin, Disperin, Primaspan, Burana, Ibusal, Ketomex, Ketorin, Orudis, Panadol, Para-tabs?

- ☐ En kertaakaan
- ☐ Harvemmin kuin 2 kertaa kuukaudessa
- ☐ Keskimäärin 2-3 kertaa kuukaudessa
- ☐ Keskimäärin 1-2 kertaa viikossa
- ☐ Keskimäärin 3-5 kertaa viikossa
- ☐ Päivittäin tai lähes päivittäin

19. Kuinka usein viimeksi kuluneen 3 kuukauden aikana olette käyttänyt vatsavaivojenne takia jotain seuraavista ripulilääkkeistä: Imocur, Imodium, Lopex, Carbo medicinalis, Tannopon?

- ☐ En kertaakaan
- ☐ Harvemmin kuin 2 kertaa kuukaudessa
- ☐ Keskimäärin 2-3 kertaa kuukaudessa
- ☐ Keskimäärin 1-2 kertaa viikossa
- ☐ Keskimäärin 3-5 kertaa viikossa
- ☐ Päivittäin tai lähes päivittäin

20. Kuinka usein viimeksi kuluneen 3 kuukauden aikana olette käyttänyt vatsavaivojenne takia jotain seuraavista ummetuksen hoitoon tarkoitetuista lääkkeistä: Pursennid, Sennapur, Metalax, Toilax, Laxoberon, Duphalac, Levolac, Loraga, Movicol, Klyx, Microlax?

- ☐ En kertaakaan
☐ Harvemmin kuin 2 kertaa kuukaudessa
☐ Keskimäärin 2-3 kertaa kuukaudessa
☐ Keskimäärin 1-2 kertaa viikossa
☐ Keskimäärin 3-5 kertaa viikossa
☐ Päivittäin tai lähes päivittäin

21. Kuinka usein viimeksi kuluneen 3 kuukauden aikana olette käyttänyt vatsavaivojenne takia jotain seuraavista kuituvalmisteista: Agiocur, Agiolax, Laxamucil, Vi-Siblin?

- ☐ En kertaakaan
☐ Harvemmin kuin 2 kertaa kuukaudessa
☐ Keskimäärin 2-3 kertaa kuukaudessa
☐ Keskimäärin 1-2 kertaa viikossa
☐ Keskimäärin 3-5 kertaa viikossa
☐ Päivittäin tai lähes päivittäin

22. Kuinka usein viimeksi kuluneen 3 kuukauden aikana olette käyttänyt vatsavaivojenne takia jotain seuraavista suolistoperäisiä kiputiloja lievittävästä reseptilääkkeistä: Buscopan, Egazil Duretter, Gastrodyn comp, Librax, Spasmo-Oxepam, Litalgin?

- ☐ En kertaakaan
☐ Harvemmin kuin 2 kertaa kuukaudessa
☐ Keskimäärin 2-3 kertaa kuukaudessa
☐ Keskimäärin 1-2 kertaa viikossa
☐ Keskimäärin 3-5 kertaa viikossa
☐ Päivittäin tai lähes päivittäin

23. Kuinka usein viimeksi kuluneen 3 kuukauden aikana olette käyttänyt vatsavaivojenne takia jotain seuraavista suoliston toimintaan vaikuttavista reseptilääkkeistä: Metopram, Primperan, Prepulsid, Doryl, Mestinon, Ubretid?

- ☐ En kertaakaan
☐ Harvemmin kuin 2 kertaa kuukaudessa
☐ Keskimäärin 2-3 kertaa kuukaudessa
☐ Keskimäärin 1-2 kertaa viikossa
☐ Keskimäärin 3-5 kertaa viikossa
☐ Päivittäin tai lähes päivittäin

24. Onko Teillä ollut viimeksi kuluneen 3 kuukauden aikana ollut käytössä jokin seuraavista masennus- eli depressiolääkkeistä:

a) Triptyl, Klotriptyl, Saroten, Pertriptyl, Suprium, Doxal, Tolvon?

- ☐ Kyllä ☐ Ei

b) Seronil, Seromex, Fontex, Fevarin, Seroxat, Cipramil?

- ☐ Kyllä ☐ Ei

c) Jokin muu masennuslääke? Mikä? (Kirjoittakaa lääkkeen kauppanimi viivalle)

25. Oletteko käyttänyt viimeksi kuluneen 3 kuukauden aikana vatsavaivojenne hoitoon joitakin muita kuin edellisissä kysymyksissä mainittuja lääkkeitä?

- ☐ En ☐ Kyllä, mitä lääkkeitä? (Kirjoittakaa lääkkeiden kauppanimet viivalle)

26. Minkä ikäinen olitte, kun vatsavaivanne alkoivat?

- ☐ Alle 20 vuotta ☐ 20-29 vuotta ☐ 30-39 vuotta ☐ 40-44 vuotta
☐ 45-49 vuotta ☐ 50-54 vuotta ☐ 55-59 vuotta ☐ 60-65 vuotta

27. Onko lähisuvussanne (vanhemmillanne, sisaruksillanne tai lapsillanne) seuraavia sairauksia:

- | | | | |
|-------------------------------|-----------------------------|--------------------------------|-----------------------------------|
| Paksu- tai peräsuolen syöpä? | <input type="checkbox"/> Ei | <input type="checkbox"/> Kyllä | <input type="checkbox"/> En tiedä |
| Haavainen paksusuolitulehdus? | <input type="checkbox"/> Ei | <input type="checkbox"/> Kyllä | <input type="checkbox"/> En tiedä |
| Crohnin tauti? | <input type="checkbox"/> Ei | <input type="checkbox"/> Kyllä | <input type="checkbox"/> En tiedä |
| Keliakia? | <input type="checkbox"/> Ei | <input type="checkbox"/> Kyllä | <input type="checkbox"/> En tiedä |

28. Onko Teillä esiintynyt viimeksi kuluneen vuoden aikana vatsavaivojenne yhteydessä seuraavia oireita:

- | | | |
|---|-----------------------------|--------------------------------|
| Verta ulosteissa? | <input type="checkbox"/> Ei | <input type="checkbox"/> Kyllä |
| Yli 3 kg laihtuminen ilman laihdutustarkoitusta? | <input type="checkbox"/> Ei | <input type="checkbox"/> Kyllä |
| Kuumeilua? | <input type="checkbox"/> Ei | <input type="checkbox"/> Kyllä |
| Herääminen yöllä vatsaoireiden takia? | <input type="checkbox"/> Ei | <input type="checkbox"/> Kyllä |
| Oireiden jatkuva paheneminen ilman lievempioireisia välikausia? | <input type="checkbox"/> Ei | <input type="checkbox"/> Kyllä |

Seuraavat kysymykset liittyvät yleiseen terveydentilaanne. Rastittakaa yksi vaihtoehto, joka parhaiten kuvaa tilannettanne.

29. Onko terveytenne yleisesti ottaen...

- ☐ erinomainen
☐ varsin hyvä
☐ hyvä
☐ tyydyttävä
☐ huono

30. Jos vertaatte nykyistä terveydentilaanne vuoden takaiseen, onko terveytenne yleisesti ottaen...

- ☐ tällä hetkellä paljon parempi kuin vuosi sitten
☐ tällä hetkellä jonkin verran parempi kuin vuosi sitten
☐ suunnilleen samanlainen
☐ tällä hetkellä jonkin verran huonompi kuin vuosi sitten
☐ tällä hetkellä paljon huonompi kuin vuosi sitten

Seuraavassa luetellaan erilaisia päivittäisiä toimintoja. Rajoittaako terveydentilaanne nykyisin suoriutumistanne seuraavista päivittäisistä toiminnoista? Jos rajoittaa, kuinka paljon? (Merkitkää joka riville yksi rasti sopivaan ruutuun)

- | | Kyllä,
rajoittaa
paljon | Kyllä,
rajoittaa
hiukan | Ei rajoita
lainkaan |
|---|-------------------------------|-------------------------------|--------------------------|
| 31. Huomattavia ponnistuksia vaativat toiminnot (esimerkiksi juokseminen, raskaiden tavaroiden nostelu, rasittava urheilu)..... | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 32. Kohtuullisia ponnistuksia vaativat toiminnot, kuten pöydän siirtäminen, imurointi, keilailu..... | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 33. Ruokakassien nostaminen tai kantaminen..... | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 34. Nouseminen portaita useita kerroksia..... | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 35. Nouseminen portaita yhden kerroksen..... | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 36. Vartalon taivuttaminen, polvistuminen, kumartuminen..... | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 37. Noin kahden kilometrin matkan kävely..... | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 38. Noin puolen kilometrin matkan kävely..... | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 39. Noin 100 metrin matkan kävely..... | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 40. Kylpeminen tai pukeutuminen..... | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |

Onko Teillä viimeisen 4 viikon aikana ollut RUUMIILLISEN TERVEYDENTILANNE TAKIA alla mainittuja ongelmia työssänne tai muissa tavanomaisissa päivittäisissä tehtävissänne? (Rastittakaa yksi ruutu joka riviltä)

	Kyllä	Ei
41. Vähensitte työhön tai muihin tehtäviin käyttämäänne aikaa	<input type="checkbox"/>	<input type="checkbox"/>
42. Saitte aikaiseksi vähemmän kuin halusitte	<input type="checkbox"/>	<input type="checkbox"/>
43. Terveystilanne asetti Teille rajoituksia joissakin työ- tai muissa tehtävissä	<input type="checkbox"/>	<input type="checkbox"/>
44. Toistänne tai tehtävistänne suoriutuminen tuotti vaikeuksia (olette joutunut esim. ponnistelemaan tavallista enemmän)	<input type="checkbox"/>	<input type="checkbox"/>

Onko Teillä viimeisen 4 viikon aikana ollut TUNNE-ELÄMÄÄN LIITTYVIEN VAIKEUKSIEN (esim. masentuneisuus tai ahdistuneisuus) takia alla mainittuja ongelmia työssänne tai muissa tavanomaisissa päivittäisissä tehtävissänne? (Rastittakaa yksi ruutu joka riviltä)

	Kyllä	Ei
45. Vähensitte työhön tai muihin tehtäviin käyttämäänne aikaa	<input type="checkbox"/>	<input type="checkbox"/>
46. Saitte aikaiseksi vähemmän kuin halusitte	<input type="checkbox"/>	<input type="checkbox"/>
47. Ette suorittanut töitänne tai muita tehtäviänne yhtä huolellisesti kuin tavallisesti	<input type="checkbox"/>	<input type="checkbox"/>

48. MISSÄ MÄÄRIN ruumiillinen terveydentilanne tai tunne-elämän vaikeudet ovat viimeisen 4 viikon aikana häirinneet tavanomaista (sosiaalista) toimintaanne perheen, ystävien, naapureiden tai muiden ihmisten parissa? (Rastittakaa yksi vaihtoehto)

- ☐ Ei lainkaan
- ☐ Hieman
- ☐ Kohtalaisesti
- ☐ Melko paljon
- ☐ Erittäin paljon

49. Kuinka voimakkaita ruumiillisia kipuja Teillä on ollut viimeisen 4 viikon aikana? (Rastittakaa yksi vaihtoehto)

- ☐ Ei lainkaan
- ☐ Hyvin lieviä
- ☐ Lieviä
- ☐ Kohtalaisia
- ☐ Voimakkaita
- ☐ Erittäin voimakkaita

50. Kuinka paljon kipu on häirinnyt tavanomaista työtänne (kotona tai kodin ulkopuolella) viimeisen 4 viikon aikana? (Rastittakaa yksi vaihtoehto)

- ☐ Ei lainkaan
- ☐ Hieman
- ☐ Kohtalaisesti
- ☐ Melko paljon
- ☐ Erittäin paljon

Seuraavat kysymykset koskevat sitä, miltä Teistä on tuntunut viimeisen 4 viikon aikana. Merkitkää kullekin riville yksi rasti siihen kohtaan, joka parhaiten kuvaa tuntemuksianne.

	koko ajan	suurimman osan aikaa	huomattavan osan aikaa	jonkin aikaa	vähän aikaa	en lainkaan
51. Tuntenut olevanne täynnä elinvoimaa.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
52. Ollut hyvin hermostunut.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
53. Tuntenut mielialanne niin matalaksi, ettei mikään ole voinut Teitä piristää.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
54. Tuntenut itsenne tyyneksi ja rauhalliseksi.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
55. Ollut täynnä tarmoa.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
56. Tuntenut itsenne alakuloiseksi ja apeaksi.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
57. Tuntenut itsenne "loppuunkuluneeksi".....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
58. Ollut onnellinen.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
59. Tuntenut itsenne väsyneeksi.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

60. KUINKA SUUREN OSAN AJASTA ruumiillinen terveydentilanne tai tunne-elämän vaikeudet ovat viimeisen 4 viikon aikana häirinneet tavanomaista sosiaalista toimintaanne (ystävien, sukulaisten, muiden ihmisten tapaaminen)? (rastittakaa yksi vaihtoehto)

- ☐ Koko ajan
☐ Suurimman osan aikaa
☐ Jonkin aikaa
☐ Vähän aikaa
☐ Ei lainkaan

Kuinka hyvin seuraavat väittämät pitävät paikkansa Teidän kohdallanne? (merkitkää yksi rasti joka riville)

	pitää ehdottomasti paikkansa	pitää enimmäkseen paikkansa	en osaa sanoa	enimmäkseen ei pidä paikkaansa	ehdottomasti ei pidä paikkaansa
61. Minusta tuntuu, että sairastun jonkin verran helpommin kuin muut ihmiset.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
62. Olen vähintään yhtä terve kuin kaikki muutkin tuntemani ihmiset.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
63. Uskon, että terveyteni tulee heikkenemään.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
64. Terveysteni on erinomainen.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Tarkistakaa vielä, että olette vastannut kaikkiin kysymyksiin ja palauttakaa kaavake oheisessa palautuskuoressa, jonka postimaksu on jo maksettu puolestanne.

KIITOS YHTEISTYÖSTÄNNE!